

REVIEW

Mechanisms and therapeutic effectiveness of pulsed electromagnetic field therapy in oncology

Maria Vadalà¹, Julio Cesar Morales-Medina², Annamaria Vallelunga³, Beniamino Palmieri¹, Carmen Laurino¹ & Tommaso Iannitti⁴

¹Department of General Surgery and Surgical Specialties, Surgical Clinic, University of Modena and Reggio Emilia Medical School, Modena, Italy

²Centro de Investigación en Reproducción Animal, CINVESTAV- Universidad Autónoma de Tlaxcala, Tlaxcala, Mexico

³Department of Medicine and Surgery, Centre for Neurodegenerative Diseases (CEMAND), University of Salerno, Salerno, Italy

⁴Department of Neuroscience, Sheffield Institute for Translational Neuroscience (SITraN), University of Sheffield, Sheffield, United Kingdom

Keywords

Cancer, electromagnetic therapy, oncology, pulsed electromagnetic fields, tumor-specific frequencies

Correspondence

Tommaso Iannitti, Sheffield Institute for Translational Neuroscience, University of Sheffield, 385A Glossop Road, Sheffield S10 2HQ, United Kingdom.
Tel: +44 7521471447;
E-mail: tommaso.iannitti@gmail.com

Funding Information

The authors thank Iskra Medical (Stegne 23, 1000 Ljubljana, Slovenia) for supporting the open access publication of this article.

Received: 23 May 2016; Revised: 18 July 2016; Accepted: 19 July 2016

Cancer Medicine 2016; 5(11):3128–3139

doi: 10.1002/cam4.861

Introduction

Cancer is one of the most common causes of death worldwide and accounted for 8.2 million deaths in 2012 [1]. The number of cancer-related deaths is predicted to increase to over 11 million by 2030 [2]. The types of cancer with the highest incidence are lung (1.59 million people), liver (745,000), stomach (723,000), colon and rectum (694,000), breast (521,000), and esophagus (400,000) [1]. In oncology, the selection of correct treatment strategy, in early disease stages, is crucial to increase the probability of remission and improve survival. Available cancer treatments include chemotherapy, immunotherapy or antibody-based therapy, radiation therapy, and surgery [3]. The therapeutic strategy is chosen taking into account the individual patient's medical assessment, type of cancer,

Abstract

Cancer is one of the most common causes of death worldwide. Available treatments are associated with numerous side effects and only a low percentage of patients achieve complete remission. Therefore, there is a strong need for new therapeutic strategies. In this regard, pulsed electromagnetic field (PEMF) therapy presents several potential advantages including non-invasiveness, safety, lack of toxicity for non-cancerous cells, and the possibility of being combined with other available therapies. Indeed, PEMF stimulation has already been used in the context of various cancer types including skin, breast, prostate, hepatocellular, lung, ovarian, pancreatic, bladder, thyroid, and colon cancer in vitro and in vivo. At present, only limited application of PEMF in cancer has been documented in humans. In this article, we review the experimental and clinical evidence of PEMF therapy discussing future perspectives in its use in oncology.

location, and disease stage [4]. Multimodal treatments are often required to reduce the therapy-induced side effects [5] related to pharmacological as well as other approaches including surgery [6]. Chemotherapy-induced side effects depend on various variables such as the drug employed, its dosage, and treatment duration. These side effects include pain, fatigue, throat and mouth sores, diarrhea, nausea, vomiting, constipation, and blood disorders. Side effects affecting the nervous system are commonly experienced with chemotherapy and include cognitive dysfunction, headache, dizziness, vision loss and vision disturbances such as blurred or double vision, changes in learning and memory, sexual dysfunction, ataxia, and peripheral neuropathy [7–11]. Rashes, fever, hypotension, colitis or other gastrointestinal problems, and thyroid dysfunctions are immunotherapy-related side effects [12]. The main

radiotherapy-induced side effects are dry mouth and gum sores, jaw stiffness, nausea, lymphedema, swallowing difficulties, shortness of breath, breast or nipple soreness, rectal bleeding, incontinence, bladder irritation, and pituitary dysfunction [13]. Surgical techniques, such as minimally invasive surgery, also result in pain, fatigue, appetite loss, swelling and bruising around the site of surgery, bleeding, infection, lymphedema, and organ dysfunction [14]. Numerous studies support the development of new treatments in oncology to be added to the traditional protocols to increase the effectiveness of available treatments, reducing side effect profile, and the patients' quality of life [15–18]. Such resources include traditional Chinese medicine, Ayurvedic medicine, homeopathy, and naturopathy [19]. While complementary and alternative medicine (CAM) is not generally considered part of conventional medicine, it has been widely used in the oncology field as an add-on therapy to control patients' symptoms and improve their quality of life [20–26]. The beginning of the 20th century saw the first therapeutic applications of CAM therapies for cancer treatment; these therapies include acupuncture, chromotherapy, therapeutic touch (reiki), and pulsed electromagnetic field (PEMF) therapy [4, 15, 27–30]. In this review, we have focused on PEMF therapy, a non-invasive technique characterized by electromagnetic fields inducing microcurrents to the entire body or locally to target specific body tissues. Exposure to PEMFs in the 0–300 Hz range is a therapeutic tool extensively used for the treatment of several pathologies including osteoarthritis, Parkinson's disease, postsurgical pain and edema, treatment of chronic wounds, and facilitation of vasodilatation and angiogenesis producing direct stimulation to excitable cells including nerve and muscle cells [31–34]. Stimulation with sufficient intensity and duration induces a current across targeted cell membranes, activating nerve cells or muscles to propagate action potentials [35–37]. Indeed, PEMF therapy can be used as an adjuvant treatment to chemotherapy and radiotherapy with the aim of reducing their dosage, mitigating any harmful secondary side effects, and enhancing patient's prognosis [15, 35, 38–40].

Aim and searching criteria

We reviewed *in vitro*, *in vivo*, and clinical studies employing PEMF therapy for cancer treatment published between 1976 and 2016. We searched Pubmed/Medline, Embase, Web of Science and Scopus using the keywords “PEMFs”, “cancer”, “magnet therapy”, “tumour specific frequencies” and “oncology” alone or combined. This review aims at describing the state of the art of PEMF therapy, discussing current understanding of the underlying mechanisms and outlining future therapeutic perspectives in oncology.

In Vitro Studies

PEMF therapy has been extensively studied *in vitro* using various human cancer cell lines, such as pheochromocytoma-derived (PC12), breast cancer (e.g., MCF7, MDA-MB-231 and T47D), and colon cancer (SW-480 and HCT-116) [41–45]. These studies have shown that PEMF therapy may exert proliferative inhibition and mitotic spindle disruption [18, 40], block the development of neovascularization required for tumor supply [46–48] and exacerbate an inherent or induced genetic instability by reducing the stringency of the late-cycle (G2) checkpoint [49]. While chemotherapy is not specific to cancer cells and targets all rapidly dividing cells [50–52], PEMFs exert selective cytotoxic effect on neoplastic cells [15, 40, 53–55] making this therapy a highly promising strategy.

In the next subparagraphs, we will review studies employing PEMF therapy in different cell lines as a model to study specific types of cancer (Table 1).

Studies of PEMF therapy in human breast cancer and colon cancer cell lines

A study by Crocetti and coworkers [38] investigated whether ultra-low intensity and frequency PEMF therapy could induce apoptosis in human breast adenocarcinoma cells (MCF7). PEMF exposure was cytotoxic to MCF7 cells, but not to normal breast epithelial cells (MCF10). Both MCF7 and MCF10 cells were exposed to PEMF therapy and the cytotoxic indices were measured in order to design PEMF paradigms that could reduce selectively neoplastic cell proliferation. The PEMF parameters tested were: (1) frequency of 20 Hz, (2) intensity of 3 mT and (3) exposure time of 60 min/day for up to 3 days. Four independent methods of monitoring cancer-induced apoptosis (trypan blue assay, apoptosis determination by DNA strand break detection, analysis of cellular electrical properties by means of impedance microflow cytometer, and apoptosis determination by Annexin V staining) showed that this specific set of PEMF parameters was cytotoxic to breast cancer cells. While this treatment selectively induced apoptosis of MCF7 cells, it had no effect on MCF10 cells that were more resistant to apoptosis in response to PEMFs. Although these results are encouraging, PEMF exposure was limited to 3 days. Long-term PEMF exposure needs to be assessed in further studies based on the concept that PEMF effectiveness is strictly linked to the signal parameters, exposure magnitude, duration, signal shape, duration of treatment as well as the type of cells exposed to the magnetic field [56, 57].

The antineoplastic effect of PEMFs has also been investigated in human breast cancer MDA-MB-231, colon cancer SW-480, and HCT-116 cell lines. These cells were exposed to 50 Hz PEMFs for 24 and 72 h [58]. PEMFs decreased the number of viable cells in all the cell lines tested,

Table 1. In vitro studies of PEMF therapy in oncology.

Author(s), year	Cell type	Treatment	Main findings	References
Crocetti et al., 2013	Human breast adenocarcinoma cells (MCF7) and nontumorigenic cells (MCF10)	Daily 60-min PEMF therapy session (20 Hz; 3 mT) for 3 days	PEMFs increased apoptosis in MCF7 cells but had no effect on MCF10 cells	[38]
Filipovic et al., 2014	Human breast cancer (MDA-MB-231) and colon cancer (SW-480 and HCT-116) cell lines	24 and 72 h exposure to PEMF therapy (50 Hz; 10 mT)	PEMFs increased apoptosis in MDA-MB-231 (55% and 20%), SW480 (11% and 6%), and HCT-116 cell lines (2% and 3%) after 24 and 72 h exposure, respectively, compared with untreated control cancer cell lines	[58]
Morabito et al., 2010	Undifferentiated PC12 pheochromocytoma cells and differentiated PC12 cells	Short PEMF therapy session (50 Hz, 0.1–1.0 mT) for 30 min, and long-term PEMF session (50 Hz, 0.1–1.0 mT) for 7 days	30-min PEMF session in undifferentiated PC12 cells increased ROS levels and decreased catalase activity. No change in intracellular Ca ²⁺ concentration was observed. 7-day PEMF therapy session in undifferentiated PC12 cells resulted in increased intracellular Ca ²⁺ concentration and increased catalase activity. No significant findings were observed in differentiated PC12 cells	[41]

PEMF, pulsed electromagnetic field; ROS, reactive oxygen species.

reaching 55% after 24 h and 20% after 72 h in the MDA-MB-231 cell line, 11% after 24 h and 6% after 72 h in the SW480 cell line, and 2% after 24 h and 3% after 72 h in the HCT-116 cell line, compared with unexposed cancer cell lines used as controls, as assessed by a computer reaction-diffusion model, a mathematical model widely employed to study cell proliferation and infiltration [59]. The lower percentage inhibition of neoplastic cell proliferation was observed after 72 h, showing that PEMF therapy had antiproliferative activity which decreased over time. This action is exerted in vitro by interfering with microtubule spindle polymerization. Indeed, PEMF exposure reduces the fraction of polymerized microtubules, disrupts the mitotic spindle structure, inhibits cell division, thereby leading to chromosome mis-segregation and cancer-induced apoptosis [60]. In summary, studies in human breast and colon cancer cell lines are promising and warrant further investigations.

Studies of PEMF therapy in pheochromocytoma-derived cells

PEMF signal parameters have been extensively utilized on diverse cell types to determine in vitro effectiveness [61, 62]. For example, Morabito and coworkers [41] investigated cell responsiveness and in vitro neuritogenesis following PEMF exposure. They specifically focused on PEMF ability to modify morphology, proliferation, and differentiation in PC12 pheochromocytoma cells. Furthermore, they assessed whether PEMFs can induce variable and species-specific

alterations in the oxidative stress pathway such as Ca²⁺-dependent oxidative stress which enhances free radical production, particularly via the Fenton reaction, leading to apoptotic cell death [63–69]. Undifferentiated and differentiated [supplemented with 50 ng/mL of nerve growth factor (NGF)] PC12 cells were exposed to 50 Hz PEMF therapy (0.1–1.0 mT), and cell growth and viability were evaluated after immediate (30 min) or long-term exposure (7 days), using colorimetric and morphological assays. The long-lasting exposure to PEMFs did not affect the biological response in terms of proliferation and neuritogenesis. Thirty-minute PEMF exposure at 1.0 mT in undifferentiated PC12 cells increased the levels of reactive oxygen species (ROS) and decreased catalase activity, an indicator of oxidative stress. Conversely, long-term PEMF exposure of undifferentiated PC12 cells also increased catalase activity that could reflect the absence of ROS accumulation and a possible adaptation cell response to PEMFs. During immediate PEMF exposure in undifferentiated PC12 cells, no change in intracellular Ca²⁺ concentration was observed, while it increased after long-term exposure. This enhanced calcium level could activate, through voltage-gated (L-type) calcium channels, signaling pathways and lead to the expression of genes modulating cell differentiation, survival, and apoptosis such as extracellular signal-regulated kinases, c-Jun N-terminal protein kinase/stress-activated protein kinase, and p38 [70–73]. In particular, the undifferentiated PC12 cells were more sensitive to PEMFs exposure, while the differentiated PC12 cells were more stable and resistant to stress,

probably due to the action of the cell surface NGF receptors such as p75NR [74].

Further studies are necessary to identify the ROS/intracellular Ca^{2+} cross-talking pathway activated by PEMF therapy. However, the study by Morabito and coworkers supports the hypothesis that ROS and Ca^{2+} could be the cellular “primum movens” of PEMF therapy-induced effects, as observed in pheochromocytoma cells.

In Vivo Studies

Several studies investigated the antineoplastic effect of PEMFs using widely employed animal models of several types of cancer, including breast cancer, hepatocellular carcinoma (HCC), and melanoma (Table 2) [4, 48, 75–78].

PEMF therapy effectiveness in mouse models of breast cancer

PEMF therapy effectiveness on tumor growth and viability has been tested in mouse models of breast cancer. For example, xenograft mouse models are widely used to study breast cancer. This model is obtained by injection of human breast cancer cells including estrogen-negative (MDA-MB-231) and estrogen-positive (MCF7) breast carcinoma cell lines or mouse breast cancer cells including EpH4 mammary epithelial cells or mitogen-activated protein kinase (MEK)-transformed EpH4 cells subcutaneously, intravenously, intracardially, or orthotopically, four times every 5 days, into the mammary fat pad of immunocompromised mice [79, 80]. The injected cells are highly invasive in vitro and tumorigenic when transplanted into the mammary fat pad. After a week from the last injection, the mouse is palpated biweekly for mammary tumors and the dimensions of tumors are measured using an external caliper daily. Mice are euthanized when the tumor size becomes ulcerated with macro-metastases, mainly in liver, bone, and brain [81–84]. For example, EpH4-MEK Bcl2¹³ cells (1×10^6) transfected with a luciferase expression vector (p β P2-PolIII-luciferase) were injected into the mammary fat pad in 12 T-cell-immunodeficient Swiss outbred female nude mice (Cr:NIH(S)-*nu/nu*) [85]. Mice were divided into four groups ($n = 3$ each). Group 1, 2, and 3 were exposed to PEMF therapy (1 Hz, 100 mT) daily for 60, 180, or 360 min, respectively, for 4 weeks, while group 4 did not receive PEMF therapy and was used as control. All mice were monitored for tumor growth by body bioluminescence imaging once every 2 to 4 days for 4 weeks. Then, all the mice were sacrificed and skin, liver, lung, and spleen samples were collected for histopathologic analysis. Mice exposed to PEMFs for 60 and 180 min daily showed a 30% and 70% breast tumor reduction, respectively, at week 4, if compared to baseline. Mice exposed to PEMFs for 360 min daily,

showed a suppression of tumor growth at week 4. In summary, this study shows that the time of PEMF exposure is critical to determine its effectiveness. Mice exposed for longer duration (360 min daily for 4 weeks) showed a significant reduction in tumor size, due probably to the inhibition of angiogenesis that may suppress the formation of blood vessels in tumor tissues, reducing the tumor growth.

Antineoplastic effect of PEMF therapy in rodent models of hepatocellular carcinoma

Chemically induced HCC is a widely used model of hepatocarcinogenesis that mimics the development of fibrosis and cirrhosis. This model is obtained by intraperitoneal administration of a carcinogenic agent, *N*-diethylnitrosamine (DEN; 50–100 mg/kg mouse body weight) alone or followed by oral administration of a nongenotoxic liver tumor promoter [phenobarbital (PB)]. DEN induces damage to DNA, proteins, and lipids, leading to hepatocyte death [86]. It is hydroxylated to α -hydroxyl nitrosamine, mediated by cytochrome P450 enzymes which are primarily located in the centrilobular hepatocytes. Then, an electrophilic ethyldiazonium ion is formed and causes DNA damage by reacting with nucleophiles. Three to four weeks following the last injection, mice receive drinking water containing PB (0.07%) that increases the expression of cytochrome P450, inducing oxidative stress and resulting in HCC development after 6 months from PB administration [86–90]. Emara and coworkers evaluated the safety and effectiveness of PEMFs with different intensity and frequency in a rat model of DEN-induced HCC (75 mg/kg body weight, once a week for 3 weeks) [91]. Sixty rats were divided into six groups: Group 1 (naive rats) received PEMF therapy (2–3 Hz, 0.004 T) for 30 min/day for 6 days/week for 4 weeks; group 2 (naive rats) received PEMF therapy (<1 Hz, 0.6 T) 15 min/day for 6 days/week for 4 weeks; group 3 (naive rats) was left untreated; group 4 (HCC rats) received PEMF therapy (2–3 Hz, 0.004 T) for 30 min/day for 6 days/week for 4 weeks; group 5 (HCC rats) received PEMF therapy (<1 Hz, 0.6 T) 15 min/day for 6 days/week for 4 weeks; group 6 (HCC rats) was left untreated. No changes in histopathology and dielectric properties of liver tissue were observed in naive rats exposed to PEMFs supporting its safety. In HCC rats exposed to PEMFs, a significant decrease in AFP level (AFP is a serum glycoprotein often elevated in HCC patients and used as a carcinoma marker in the clinic) was reported together with a slight improvement in dielectric properties of liver tissue. These results were confirmed by electron microscopy and histological analysis showing HCC regression. Altogether this evidence supports the antineoplastic activity of PEMF therapy in

Table 2. In vivo studies of PEMF therapy in oncology.

Author(s), year	Animal model (number of animals, study design)	Route of administration	Treatment	Main findings	References
Tatarov et al., 2011	12 T-cell-immunodeficient Swiss outbred female nude mice (Cr:NIH(S)- <i>nu/nu</i>), divided into 4 groups ($n = 3$ each)	Orthotopic injection of metastatic mouse breast tumor cell line [Eph4-MEK Bcl2 ¹³ cells (1×10^6)] into the mammary fat pad	Group 1, 2 and 3 were exposed to PEMFs (1 Hz, 100 mT) daily for 60, 180, or 360 min, respectively, for 4 weeks; group 4 did not receive any treatment and was used as control	Mice exposed for 60 and 180 min daily showed a 30% and 70% tumor reduction, respectively, at week 4, if compared to baseline	[85]
Emara et al., 2013	60 rats (strain not reported) divided into 6 groups	Intraperitoneal administration of a carcinogenic agent, DEN	Group 1 (naive rats) received PEMF therapy (2-3 Hz, 0.004 T) for 30 min/day for 6 days/week for 4 weeks; group 2 (naive rats) received PEMF therapy (<1 Hz, 0.6 T) 15 min/day for 6 days/week for 4 weeks; group 3 (naive rats) was left untreated; group 4 (HCC rats) received PEMF therapy (2-3 Hz, 0.004 T) for 30 min/day for 6 days/week for 4 weeks; group 5 (HCC rats) received PEMF therapy (<1 Hz, 0.6 T) 15 min/day for 6 days/week for 4 weeks; group 6 (HCC rats) was left untreated.	A significant decrease in serum AFP level and a slight improvement in dielectric properties of liver tissues was observed in HCC rats treated with PEMFs. These results were confirmed by electron microscopy and histological analysis showing HCC regression. No changes in histopathology and dielectric properties of liver tissue were observed in naive rats exposed to PEMFs.	[91]
Nuccitelli et al., 2006	23 SKH-1 immunocompetent, hairless, albino mice	Single subcutaneous injection of B16 murine melanoma cells (1×10^5) on the dorsal side of the mouse ear	30-min PEMF therapy session (0.5 Hz, 0.2 T) three times a day for 6 days	All mice exhibited significant pyknosis, shrinkage of the tumor cell nuclei by 54% within a few minutes after PEMF therapy and by 68% within 3 h and reduction in the blood flow in about 15 min following PEMF therapy	[95]
Nuccitelli et al., 2010	Four female immunodeficient, hairless, albino Nu/Nu mice	Single subcutaneous injection of murine melanoma cells (B16-F10-eGFP, 1×10^5) on the mouse skin	Daily 6-min PEMF session (5–7 Hz, 0.2 T) for 10 days	Melanoma cells shrank within an hour post PEMF therapy, exhibiting pyknosis within 24 h post treatment. PEMFs-treated mice showed complete remission of melanoma	[100]

PEMF, pulsed electromagnetic field; DEN, *N*-diethylnitrosamine; AFP, alpha-fetoprotein; HCC, hepatocellular carcinoma

the rat model of DEN-induced HCC and warrants further investigations.

PEMF therapy effectiveness in murine melanoma models

The most frequently used murine melanoma model is the syngeneic B16 model. It is obtained by a single subcutaneous injection of 1×10^5 B16 murine melanoma cells on

the dorsal side of the mouse ear. Melanoma nodules 5–6 mm in diameter develop 7 days post-injection [92–94]. The melanoma model in SKH-1 hairless mice has been used to investigate the effectiveness of PEMF therapy (0.5 Hz, 0.2 T, 30 min/day). Mice ($n = 23$) received 1–3 PEMF treatments daily for 6 days and were monitored for tumor growth, daily, by optical methods, such as transillumination and power Doppler ultrasound reconstructions that display blood flow images for each tumor [95].

Then, all the mice were sacrificed and skin tissues were collected for histopathological analysis. All mice exposed to PEMFs exhibited significant pyknosis, shrinkage of the tumor cell nuclei by 54% within a few minutes after PEMF therapy and by 68% within 3 h and reduction in the blood flow in about 15 min following PEMF therapy. These effects may be due to PEMF therapy that stimulates murine melanoma to self-destruct by triggering rapid pyknosis of tumor cell nuclei and reducing blood flow [96–99]. A further study [100] optimized the PEMF therapy parameters pulse number, amplitude, and frequency to completely suppress melanoma with a single treatment. In this study, four female immunodeficient, hairless, albino Nu/Nu mice received a single PEMF treatment for 6 min using the following parameters: 2.700 pulses, amplitude of 30 kV/cm and frequency of 5–7 Hz for 10 days. After 2–4 weeks, mice were sacrificed and skin samples were processed for histology. Melanoma cells shrank within an hour post PEMF therapy, exhibiting pyknosis within 24 h post PEMFs and showing a complete remission of melanoma in all the mice, as assessed by *in vivo* imaging (transillumination and photography). To evaluate the safety of PEMF therapy, the authors recorded the physiological parameters and introduced a miniature thermocouple into the tumor for simultaneous measurement of intratumoral temperature during PEMF treatment; body temperature and systolic blood pressure showed no significant changes, while the intratumoral temperature was ~6–7°C, evidencing that, by limiting the frequency to 7 Hz or less, it was possible to avoid heating the tumor to hyperthermia temperatures potentially leading to damage of the surrounding tissues. Evidence of efficacy of a single PEMF treatment on mouse skin cancer resulting in suppression of tumor growth and induction of apoptosis is promising for translational applications.

Clinical Studies

The use of PEMF therapy in oncology is still limited (Table 3) [4]. The first study utilizing PEMF therapy was conducted by Barbault and coworkers who hypothesized that a combination of specific frequencies, defined tumor-specific frequencies, may display therapeutic effectiveness for localized treatment of tumors [15]. They identified a total of 1524 tumor-specific frequencies, ranging from 0.1 to 114 kHz, consisting in the measurement of variations in skin electrical resistance, pulse amplitude, and blood pressure in 163 patients affected by different types of cancer including brain tumors, colorectal cancer, HCC carcinoma, pancreatic, colorectal, ovarian, breast, prostate, lung, thyroid, and bladder cancer and exposed to the radiofrequency system. Self-administered PEMF therapy for 60 min, three times a day, for an average of 278.4 months was offered

to only 28 patients with advanced cancer (breast cancer [$n = 7$], ovarian cancer [$n = 5$], pancreatic cancer [$n = 3$], colorectal cancer [$n = 2$], prostate cancer [$n = 2$], glioblastoma multiforme [$n = 1$], HCC carcinoma [$n = 1$], mesothelioma [$n = 1$], neuroendocrine tumor [$n = 1$], non-small-cell lung cancer [$n = 1$], oligodendroglioma [$n = 1$], small-cell lung cancer [$n = 1$], sarcoma [$n = 1$] and thyroid tumor [$n = 1$]). None of the patients who received PEMF therapy reported any side effects; four patients presented stable disease for 3 years (thyroid cancer with biopsy-proven lung metastases), 6 months (mesothelioma metastatic to the abdomen), 5 months (non-small-cell lung cancer), and 4 months (pancreatic cancer with biopsy-proven liver metastases), respectively.

PEMF therapy has also been employed for the treatment of HCC. Therapies for this disease are needed, especially for patients at an advanced disease stage who cannot tolerate chemotherapy or intrahepatic interventions because of impaired liver function [101]. The feasibility of PEMF therapy for treatment of HCC has also been investigated in a single-group, open-label, phase I/II clinical study [102]. Forty-one patients with advanced HCC received very low levels of PEMFs modulated at HCC-specific frequencies (100 Hz–21 kHz) and received three-daily 60 min outpatient treatments. No adverse reactions were observed during PEMF treatment. Five patients reported complete disappearance and two patients reported decrease in pain shortly after beginning of treatment. Four patients showed a partial response to treatment, while 16 patients (39%) had stable disease for more than 12 weeks. This study shows that PEMF therapy provides a safe and well-tolerated treatment, as well as evidence of antineoplastic effects in patients with HCC.

In summary, encouraging findings warrant randomized clinical studies to determine the effectiveness of amplitude-modulated PEMF therapy that can delay cancer progression and increase overall survival in patients. The increased knowledge of tumor-specific frequencies and the preliminary evidence that additional tumor-specific frequencies may yield a therapeutic benefit provide a strong rationale for the novel concept that administration of a large number of these frequencies may result in successful long-term disease management.

Discussion and Conclusions

In vitro studies support antineoplastic and antiangiogenic effects of PEMF therapy. Several mechanisms of PEMF therapy have been elucidated. For example, PEMFs inhibit cancer growth by disrupting the mitotic spindle in a process mediated by interference of spindle tubulin orientation and induction of dielectrophoresis. Furthermore, PEMF therapy modulates gene expression and protein synthesis

Table 3. Clinical studies of PEMF therapy in oncology.

Author(s), year	Study design	Number of patients	Pathology	Treatment	Outcomes	Side effects	References
Barbault et al., 2009	Compassionate and investigational clinical trial	28	Glioblastoma multiforme, mesothelioma, oligodendroglioma, sarcoma, HCC and breast, colorectal, lung, neuroendocrine, ovarian, pancreatic, prostate and thyroid cancers	60-min PEMF session (0.1 Hz–114 kHz, 1.5 T) three times a day for 278.4 months	One patient with thyroid cancer, one patient with mesothelioma metastatic to the abdomen, one patient with non-small-cell lung cancer and one patient with pancreatic cancer with biopsy-proven liver metastases presented stable disease for 3 years, 6 months, 5 months and 4 months, respectively	None reported	[15]
Costa et al., 2007	A single-group, open-label, phase I/II clinical trial	41	Advanced HCC	Daily 60-min PEMF session (100 Hz–21 kHz, 1.5 T) three times a day for 6 months	Five patients reported complete disappearance and two patients reported decrease in pain shortly after treatment. Four patients showed a partial response to treatment, while 16 patients had stable disease for more than 12 weeks	None reported	[102]

PEMF, pulsed electromagnetic field; HCC, hepatocellular carcinoma.

interacting with specific DNA sequences within gene promoter regions [18, 38, 40, 41, 58, 103]. In addition, PEMFs inhibit angiogenesis in tumor tissues, suppressing tumor vascularization and reducing tumor growth, as shown by *in vivo* studies [95–99, 104].

The specific claim, supported by the described *in vivo* studies, is that all treated groups showed slower tumor growth rate if compared with untreated control group, confirming that PEMF therapy can modulate the physiology and electrochemistry of cancer cells and influence cell membrane systems and mitosis. In addition, PEMFs induce some changes in membrane transport capacity through impacting the osmotic potential, ionic valves and leading to reduction in cellular stress factors, increase in the rate of DNA transcription, and modulation of immune response [105].

PEMFs have also an immunomodulatory effect, as supported by *in vivo* evidence showing an increase in tumor necrosis factor alpha levels that induce an anti-tumoral response, leading to the activation of a proapoptotic pathway induced by caspase-8 interaction with Fas-associated death domain, in the spleen of the murine melanoma mouse model after a 16-day therapy [78]. Changes in blood pressure, skin electrical resistance, and pulse amplitude in 163 oncology patients exposed to tumor-specific PEMF frequencies have also been reported suggesting that PEMF therapy does not only target neoplastic cells, but may also have systemic effects [15]. However, long-term PEMF treatment in HCC patients is not toxic, confirming the safety of PEMF therapy that employs 100,000 times lower frequencies if compared with radiofrequency ablation that is also employed for treatment of HCC [55].

In conclusion, only two clinical studies have used PEMF therapy for cancer treatment. These studies show that PEMF therapy is safe and promising compared to other available cancer therapies. In the future, PEMFs could be used not only as primary therapy but also in combination with other common antineoplastic therapies. Given that new portable and affordable PEMF devices are increasingly available on the market, future controlled clinical studies are expected to further determine the potential of PEMF therapy in oncology.

Acknowledgments

JCMM acknowledges CONACyT for membership.

Conflict of Interest

The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

References

- Artega, C. L., P. C. Adamson, J. A. Engelman, M. Foti, R. B. Gaynor, S. G. Hilsenbeck, et al. 2014. AACR cancer progress report 2014. *Clin. Cancer Res.* 20(19 Suppl.):S1–S112.
- Woods, R. R. C. M., and A. J. Coldman. 2015. Cancer incidence in British Columbia expected to grow by 57% from 2012 to 2030. *B. C. Med. J.* 5:190–196.
- Fischer, G., R. B. Pelka, and J. Barovic. 2005. [Adjuvant treatment of knee osteoarthritis with weak pulsing magnetic fields. Results of a placebo-controlled trial prospective clinical trial]. *Z. Orthop. Ihre Grenzgeb.* 143:544–550.
- Verginadis, I. V. A., I. Karagounis, Y. Simos, D. Peschos, S. Karkabounas, and A. Evangelou. 2012. Beneficial effects of electromagnetic radiation in Cancer. *Electromagnetic Radiation*; in tech.
- Wang, Z. M., G. X. Wang, S. M. Wang, A. Bassi, M. Khalid, K. H. Schoenbach, et al. 2006. Early response genes to DNA damage by nanosecond pulsed electric fields. *IEEE Trans. Plasma Sci.* 28:206–223.
- Tofani, S., D. Barone, M. Cintorino, M. M. de Santi, A. Ferrara, R. Orlassino, et al. 2001. Static and ELF magnetic fields induce tumor growth inhibition and apoptosis. *Bioelectromagnetics* 22:419–428.
- Coates, A., S. Abraham, S. B. Kaye, T. Sowerbutts, C. Frewin, R. M. Fox, et al. 1983. On the receiving end—patient perception of the side-effects of cancer chemotherapy. *Eur. J. Cancer Clin. Oncol.* 19:203–208.
- Schagen, S. B., M. J. Muller, W. Boogerd, G. J. Mellenbergh, and F. S. van Dam. 2006. Change in cognitive function after chemotherapy: a prospective longitudinal study in breast cancer patients. *J. Natl. Cancer Inst.* 98:1742–1745.
- Winocur, G., J. Vardy, M. A. Binns, L. Kerr, and I. Tannock. 2006. The effects of the anti-cancer drugs, methotrexate and 5-fluorouracil, on cognitive function in mice. *Pharmacol. Biochem. Behav.* 85:66–75.
- Reiriz, A. B., G. K. Reolon, T. Preissler, J. O. Rosado, J. A. Henriques, R. Roesler, et al. 2006. Cancer chemotherapy and cognitive function in rodent models: memory impairment induced by cyclophosphamide in mice. *Clin. Cancer Res.* 12:5000; author reply 00-1.
- Joshi, G., R. Sultana, J. Tangpong, M. P. Cole, D. K. St Clair, M. Vore, et al. 2005. Free radical mediated oxidative stress and toxic side effects in brain induced by the anti cancer drug adriamycin: insight into chemobrain. *Free Radic. Res.* 39:1147–1154.
- Belldegrun, A., D. E. Webb, H. A. Austin III, S. M. Steinberg, D. E. White, W. M. Linehan, et al. 1987. Effects of interleukin-2 on renal function in patients receiving immunotherapy for advanced cancer. *Ann. Intern. Med.* 106:817–822.
- al-Mefty, O., J. E. Kersh, A. Routh, and R. R. Smith. 1990. The long-term side effects of radiation therapy for benign brain tumors in adults. *J. Neurosurg.* 73:502–512.
- Stege, R. 2000. Potential side-effects of endocrine treatment of long duration in prostate cancer. *Prostate Suppl.* 10:38–42.
- Barbault, A., F. P. Costa, B. Bottger, R. F. Munden, F. Bomholt, N. Kuster, et al. 2009. Amplitude-modulated electromagnetic fields for the treatment of cancer: discovery of tumor-specific frequencies and assessment of a novel therapeutic approach. *J. Exp. Clin. Cancer Res.* 28:51.
- Evangelou, A., I. Toliopoulos, C. Giotis, A. Metsios, I. Verginadis, Y. Simos, et al. 2011. Functionality of natural killer cells from end-stage cancer patients exposed to resonant electromagnetic fields. *Electromagn. Biol. Med.* 30:46–56.
- Ronchetto, F., D. Barone, M. Cintorino, M. Berardelli, S. Lissolo, R. Orlassino, et al. 2004. Extremely low frequency-modulated static magnetic fields to treat cancer: a pilot study on patients with advanced neoplasm to assess safety and acute toxicity. *Bioelectromagnetics* 25:563–571.
- Kirson, E. D., V. Dbaly, F. Tovarys, J. Vymazal, J. F. Soustiel, A. Itzhaki, et al. 2007. Alternating electric fields arrest cell proliferation in animal tumor models and human brain tumors. *Proc. Natl. Acad. Sci. USA* 104:10152–10157.
- Committee on the Use of Complementary and Alternative Medicine by the American Public. 2005. Pp. 1–360. *Complementary and Alternative Medicine in the United States*. National Academies Press.

20. Weiger, W. A., M. Smith, H. Boon, M. A. Richardson, T. J. Kaptchuk, and D. M. Eisenberg. 2002. Advising patients who seek complementary and alternative medical therapies for cancer. *Ann. Intern. Med.* 137:889–903.
21. Adams, J., D. W. Sibbritt, G. Easthope, and A. F. Young. 2003. The profile of women who consult alternative health practitioners in Australia. *Med. J. Aust.* 179:297–300.
22. Chrystal, K., S. Allan, G. Forgeson, and R. Isaacs. 2003. The use of complementary/alternative medicine by cancer patients in a New Zealand regional cancer treatment centre. *N. Z. Med. J.* 116:U296.
23. Lee, M. M., J. S. Chang, B. Jacobs, and M. R. Wrensch. 2002. Complementary and alternative medicine use among men with prostate cancer in 4 ethnic populations. *Am. J. Public Health* 92:1606–1609.
24. Eisenberg, D. M., R. B. Davis, S. L. Ettner, S. Appel, S. Wilkey, M. Van Rompay, et al. 1998. Trends in alternative medicine use in the United States, 1990–1997: results of a follow-up national survey. *JAMA* 280:1569–1575.
25. Chang, K. H., R. Brodie, M. A. Choong, K. J. Sweeney, and M. Kerin. 2011. Complementary and alternative medicine use in oncology: a questionnaire survey of patients and health care professionals. *BMC Cancer* 11:196.
26. Barrie, R. C., and G. Deng. 2004. Complementary and alternative therapies for cancer. *Oncologist* 9:80–89.
27. Zimmerman, J. W., H. Jimenez, M. J. Pennison, I. Brezovich, D. Morgan, A. Mudry, et al. 2013. Targeted treatment of cancer with radiofrequency electromagnetic fields amplitude-modulated at tumor-specific frequencies. *Chin. J. Cancer* 32:573–581.
28. Lu, W., E. Dean-Clower, A. Doherty-Gilman, and D. S. Rosenthal. 2008. The value of acupuncture in cancer care. *Hematol. Oncol. Clin. North Am.* 22:631–648, viii.
29. Azeemi, S. T., and S. M. Raza. 2005. A critical analysis of chromotherapy and its scientific evolution. *Evid. Based Complement. Alternat. Med.* 2:481–488.
30. Miles, P. 2007. Reiki for mind, body, and spirit support of cancer patients. *Adv. Mind Body Med.* 22:20–26.
31. Iannitti, T., G. Fistetto, A. Esposito, V. Rottigni, and B. Palmieri. 2013. Pulsed electromagnetic field therapy for management of osteoarthritis-related pain, stiffness and physical function: clinical experience in the elderly. *Clin. Interv. Aging* 8:1289–1293.
32. Vadalà, M., A. Vallelunga, L. Palmieri, B. Palmieri, J. C. Morales-Medina, and T. Iannitti. 2015. Mechanisms and therapeutic applications of electromagnetic therapy in Parkinson's disease. *Behav. Brain Funct.* 11:26.
33. Ryang We, S., Y. H. Koog, K. I. Jeong, and H. Wi. 2013. Effects of pulsed electromagnetic field on knee osteoarthritis: a systematic review. *Rheumatology (Oxford)* 52:815–824.
34. Strauch, B., C. Herman, R. Dabb, L. J. Ignarro, and A. A. Pilla. 2009. Evidence-based use of pulsed electromagnetic field therapy in clinical plastic surgery. *Aesthet. Surg. J.* 29:135–143.
35. Elson, E. I. 2009. The little explored efficacy of magnetic fields in cancer treatment and postulation of the mechanism of action. *Electromagn. Biol. Med.* 28:275–282.
36. Repacholi, M. H., and B. Greenebaum. 1999. Interaction of static and extremely low frequency electric and magnetic fields with living systems: health effects and research needs. *Bioelectromagnetics* 20:133–160.
37. Organization WH. Electromagnetic fields and public health exposure to extremely low frequency fields. 2012.
38. Crocetti, S., C. Beyer, G. Schade, M. Egli, J. Frohlich, and A. Franco-Obregon. 2013. Low intensity and frequency pulsed electromagnetic fields selectively impair breast cancer cell viability. *PLoS One* 8:e72944.
39. Cameron, I. L., L. Z. Sun, N. Short, W. E. Hardman, and C. D. Williams. 2005. Therapeutic electromagnetic field (TEMF) and gamma irradiation on human breast cancer xenograft growth, angiogenesis and metastasis. *Cancer Cell Int.* 5:23.
40. Zimmerman, J. W., M. J. Pennison, I. Brezovich, N. Yi, C. T. Yang, R. Ramaker, et al. 2012. Cancer cell proliferation is inhibited by specific modulation frequencies. *Br. J. Cancer* 106:307–313.
41. Morabito, C., S. Guarnieri, G. Fano, and M. A. Mariggio. 2010. Effects of acute and chronic low frequency electromagnetic field exposure on PC12 cells during neuronal differentiation. *Cell. Physiol. Biochem.* 26:947–958.
42. Chen, Y. C., C. C. Chen, W. Tu, Y. T. Cheng, and F. G. Tseng. 2010. Design and fabrication of a microplatform for the proximity effect study of localized ELF-EMF on the growth of in vitro HeLa and PC-12 cells. *J. Micromech. Microeng.* 20:12.
43. Tuffet, S., R. de Seze, J. M. Moreau, and B. Veyret. 1993. Effects of a strong pulsed magnetic field on the proliferation of tumour cells in vitro. *Bioelectrochem. Bioenerg.* 30:151–160.
44. Sadeghipour, R., S. Ahmadian, B. Bolouri, Y. Pazhang, and M. Shafiezadeh. 2012. Effects of extremely low-frequency pulsed electromagnetic fields on morphological and biochemical properties of human breast carcinoma cells (T47D). *Electromagn. Biol. Med.* 31:425–435.
45. Loja, T., O. Stehlikova, L. Palko, K. Vrba, I. Rampl, and M. Klabusay. 2014. Influence of pulsed electromagnetic and pulsed vector magnetic potential field on the growth of tumor cells. *Electromagn. Biol. Med.* 33:190–197.

46. Fang, M., H. Zhang, and S. Xue. 1998. Role of calcium in apoptosis of HL-60 cells induced by harringtonine. *Sci. China C Life Sci.* 41:600–607.
47. da Silva, C. P., C. R. de Oliveira, M. da Conceicao, and P. de Lima. 1996. Apoptosis as a mechanism of cell death induced by different chemotherapeutic drugs in human leukemic T-lymphocytes. *Biochem. Pharmacol.* 51:1331–1340.
48. Zhang, X., H. Zhang, C. Zheng, C. Li, and W. Xiong. 2002. Extremely low frequency (ELF) pulsed-gradient magnetic fields inhibit malignant tumour growth at different biological levels. *Cell Biol. Int.* 26:599–603.
49. Harris, P. A., J. Lamb, B. Heaton, and D. N. Wheatley. 2002. Possible attenuation of the G2 DNA damage cell cycle checkpoint in HeLa cells by extremely low frequency (ELF) electromagnetic fields. *Cancer Cell Int.* 2:3.
50. DeHaven, C. 2014. Chemotherapy and radiotherapy effects on the skin. *Plast. Surg. Nurs.* 34:192–195.
51. Corcos, D. 2013. Toward a universal treatment for cancer: cell inflation assisted chemotherapy. *Cancer Med.* 2:421–426.
52. Barton-Burke, M., G. M. Wilkes, and K. Ingwersen. 2001. P. 778 *in* Cancer chemotherapy: a nursing process approach. Jones & Bartlett Learning Publishers.
53. Barton-Burke, M., and G. M. Wilkes. 2004. Pp. 1–376 *in* Cancer therapies. Jones & Bartlett Learning Publishers.
54. Crocetti, S., F. Piantelli, and C. Leonzio. 2011. Selective destabilization of tumor cells with pulsed electric and magnetic sequences: a preliminary report. *Electromagn. Biol. Med.* 30:128–135.
55. Costa, F. P., A. C. de Oliveira, R. Meirelles, M. C. Machado, T. Zanesco, R. Surjan, et al. 2011. Treatment of advanced hepatocellular carcinoma with very low levels of amplitude-modulated electromagnetic fields. *Br. J. Cancer* 105:640–648.
56. Radeva, M., and H. Berg. 2004. Differences in lethality between cancer cells and human lymphocytes caused by LF-electromagnetic fields. *Bioelectromagnetics* 25:503–507.
57. Koh, E. K., B. K. Ryu, D. Y. Jeong, I. S. Bang, M. H. Nam, and K. S. Chae. 2008. A 60-Hz sinusoidal magnetic field induces apoptosis of prostate cancer cells through reactive oxygen species. *Int. J. Radiat. Biol.* 84:945–955.
58. Filipovic, N. D. T., M. Radovic, D. Cvetkovic, M. Curcic, S. Markovic, A. Peulic, et al. 2014. Electromagnetic field investigation on different cancer cell lines. *Cancer Cell Int.* 14:1–10.
59. Swanson, K. R., C. Bridge, J. D. Murray, and E. C. Alvord Jr.. 2003. Virtual and real brain tumors: using mathematical modeling to quantify glioma growth and invasion. *J. Neurol. Sci.* 216:1–10.
60. Filipovic, N. D., A. S. Peulic, N. D. Zdravkovic, V. M. Grbovic-Markovic, and A. J. Jurisic-Skevin. 2011. Transient finite element modeling of functional electrical stimulation. *Gen. Physiol. Biophys.* 30:59–65.
61. Simko, M. 2007. Cell type specific redox status is responsible for diverse electromagnetic field effects. *Curr. Med. Chem.* 14:1141–1152.
62. Coskun, S., B. Balabanli, A. Canseven, and N. Seyhan. 2009. Effects of continuous and intermittent magnetic fields on oxidative parameters in vivo. *Neurochem. Res.* 34:238–243.
63. Grassi, C., M. D'Ascenzo, A. Torsello, G. Martinotti, F. Wolf, A. Cittadini, et al. 2004. Effects of 50 Hz electromagnetic fields on voltage-gated Ca²⁺ channels and their role in modulation of neuroendocrine cell proliferation and death. *Cell Calcium* 35:307–315.
64. Liboff, A. R., S. Cherng, K. A. Jenrow, and A. Bull. 2003. Calmodulin-dependent cyclic nucleotide phosphodiesterase activity is altered by 20 microT magnetostatic fields. *Bioelectromagnetics* 24:32–38.
65. Manikonda, P. K., P. Rajendra, D. Devendranath, B. Gunasekaran, B. Channakeshava, R. S. Aradhya, et al. 2007. Influence of extremely low frequency magnetic fields on Ca²⁺ signaling and NMDA receptor functions in rat hippocampus. *Neurosci. Lett.* 413:145–149.
66. Morabito, C., F. Rovetta, M. Bizzarri, G. Mazzoleni, G. Fano, and M. A. Mariggio. 2010. Modulation of redox status and calcium handling by extremely low frequency electromagnetic fields in C2C12 muscle cells: a real-time, single-cell approach. *Free Radic. Biol. Med.* 48:579–589.
67. Vyklicky, L., Jr.. 1993. Calcium-mediated modulation of N-methyl-D-aspartate (NMDA) responses in cultured rat hippocampal neurons. *J. Physiol.* 470:575–600.
68. Mariggio, M. A., C. Morabito, G. Fanò Illic', F. Cuccurullo, C. Di Ilio. 2006. Extremely low frequency electromagnetic fields and oxidative stress in excitable cell lines. *Biol. Eff. Electromagn. Fields* 4(1):1043–1050.
69. Barrera-Garcia, A., T. O'Hara, F. Galvan-Magana, L. C. Mendez-Rodriguez, J. M. Castellini, and T. Zenteno-Savin. 2013. Trace elements and oxidative stress indicators in the liver and kidney of the blue shark (*Prionace glauca*). *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 165:483–490.
70. Nicotera, P., and S. Orrenius. 1998. The role of calcium in apoptosis. *Cell Calcium* 23:173–180.
71. Bito, H., K. Deisseroth, and R. W. Tsien. 1997. Ca²⁺-dependent regulation in neuronal gene expression. *Curr. Opin. Neurobiol.* 7:419–429.
72. Morgado-Valle, C., L. Verdugo-Diaz, D. E. Garcia, C. Morales-Orozco, and R. Drucker-Colin. 1998. The role of voltage-gated Ca²⁺ channels in neurite growth of cultured chromaffin cells induced by extremely low

- frequency (ELF) magnetic field stimulation. *Cell Tissue Res.* 291:217–230.
73. Dolmetsch, R. E., U. Pajvani, K. Fife, J. M. Spotts, and M. E. Greenberg. 2001. Signaling to the nucleus by an L-type calcium channel-calmodulin complex through the MAP kinase pathway. *Science* 294:333–339.
 74. Ingraham, C. A., and N. F. Schor. 2009. Necdin and TrkA contribute to modulation by p75NTR of resistance to oxidant stress. *Exp. Cell Res.* 315:3532–3542.
 75. Wang, Z., P. Yang, H. Xu, A. Qian, L. Hu, and P. Shang. 2009. Inhibitory effects of a gradient static magnetic field on normal angiogenesis. *Bioelectromagnetics* 30:446–453.
 76. Tofani, S., M. Cintonino, D. Barone, M. Berardelli, M. M. De Santi, A. Ferrara, et al. 2002. Increased mouse survival, tumor growth inhibition and decreased immunoreactive p53 after exposure to magnetic fields. *Bioelectromagnetics* 23:230–238.
 77. Williams, C. D., M. S. Markov, W. E. Hardman, and I. L. Cameron. 2001. Therapeutic electromagnetic field effects on angiogenesis and tumor growth. *Anticancer Res.* 21(6A):3887–3891.
 78. Yamaguchi, S., M. Ogiue-Ikeda, M. Sekino, and S. Ueno. 2006. Effects of pulsed magnetic stimulation on tumor development and immune functions in mice. *Bioelectromagnetics* 27:64–72.
 79. Tofani, S., D. Barone, S. Peano, P. Ossola, F. Ronchetto, and M. Cintonino. 2002. Anticancer activity by magnetic fields: inhibition of metastatic spread and growth in a breast cancer model. *Trans. Plasma Sci.* 30:1552–1557.
 80. Berg, H., B. Gunther, I. Hilger, M. Radeva, N. Traitcheva, and L. Wollweber. 2010. Bioelectromagnetic field effects on cancer cells and mice tumors. *Electromagn. Biol. Med.* 29:132–143.
 81. Pinkas, J., and P. Leder. 2002. MEK1 signaling mediates transformation and metastasis of EpH4 mammary epithelial cells independent of an epithelial to mesenchymal transition. *Cancer Res.* 62:4781–4790.
 82. Reichmann, E., H. Schwarz, E. M. Deiner, I. Leitner, M. Eilers, J. Berger, et al. 1992. Activation of an inducible c-FosER fusion protein causes loss of epithelial polarity and triggers epithelial-fibroblastoid cell conversion. *Cell* 71:1103–1116.
 83. Pinkas, J., S. S. Martin, and P. Leder. 2004. Bcl-2-mediated cell survival promotes metastasis of EpH4 betaMEKDD mammary epithelial cells. *Mol. Cancer Res.* 2:551–556.
 84. Iorns, E., K. Drews-Elger, T. M. Ward, S. Dean, J. Clarke, D. Berry, et al. 2012. A new mouse model for the study of human breast cancer metastasis. *PLoS One* 7:e47995.
 85. Tatarov, I., A. Panda, D. Petkov, K. Kolappaswamy, K. Thompson, A. Kavirayani, et al. 2011. Effect of magnetic fields on tumor growth and viability. *Comp. Med.* 61:339–345.
 86. Heindryckx, F., I. Colle, and H. Van Vlierberghe. 2009. Experimental mouse models for hepatocellular carcinoma research. *Int. J. Exp. Pathol.* 90:367–386.
 87. Goldfarb, S., T. D. Pugh, H. Koen, and Y. Z. He. 1983. Preneoplastic and neoplastic progression during hepatocarcinogenesis in mice injected with diethylnitrosamine in infancy. *Environ. Health Perspect.* 50:149–161.
 88. Koen, H., T. D. Pugh, and S. Goldfarb. 1983. Centrilobular distribution of diethylnitrosamine-induced hepatocellular foci in the mouse. *Lab. Invest.* 49:78–81.
 89. Newell, P., A. Villanueva, S. L. Friedman, K. Koike, and J. M. Llovet. 2008. Experimental models of hepatocellular carcinoma. *J. Hepatol.* 48:858–879.
 90. Leenders, M. W., M. W. Nijkamp, and I. H. Borel Rinke. 2008. Mouse models in liver cancer research: a review of current literature. *World J. Gastroenterol.* 14:6915–6923.
 91. Emara, S. O., S. M. EL-Kholy, A. H. Kazem, N. A. Hussein, and R. S. Shams Al-dein. 2013. Therapeutic effects of low frequency pulsed electromagnetic fields on rat liver cancer. *Res. Inventy Int. J. Eng. Sci.* 2:17–18.
 92. Overwijk, W. W., and N. P. Restifo. 2001. B16 as a mouse model for human melanoma. *Curr. Protoc. Immunol.* Chapter 20:Unit 20.1.
 93. Chen, X., R. James Swanson, J. F. Kolb, R. Nuccitelli, and K. H. Schoenbach. 2009. Histopathology of normal skin and melanomas after nanosecond pulsed electric field treatment. *Melanoma Res.* 19:361–371.
 94. Kranjc, S., M. Kranjc, J. Scancar, J. Jelenc, G. Sersa, and D. Miklavcic. 2016. Electrochemotherapy by pulsed electromagnetic field treatment (PEMF) in mouse melanoma B16F10 in vivo. *Radiol. Oncol.* 50:39–48.
 95. Nuccitelli, R., U. Pliquett, X. Chen, W. Ford, R. James Swanson, S. J. Beebe, et al. 2006. Nanosecond pulsed electric fields cause melanomas to self-destruct. *Biochem. Biophys. Res. Commun.* 343:351–360.
 96. White, J. A., P. F. Blackmore, K. H. Schoenbach, and S. J. Beebe. 2004. Stimulation of capacitative calcium entry in HL-60 cells by nanosecond pulsed electric fields. *J. Biol. Chem.* 279:22964–22972.
 97. Beebe, S. J., P. Fox, L. J. Rec, K. Somers, R. H. Stark, and K. H. Schoenbach. 2002. Nanosecond pulsed electric field (nsPEF) effects on cells and tissues: apoptosis induction and tumor growth inhibition. *IEEE Trans. Plasma Sci.* 30:286–292.
 98. Beebe, S. J., P. F. Blackmore, J. White, R. P. Joshi, and K. H. Schoenbach. 2004. Nanosecond pulsed electric fields modulate cell function through intracellular signal transduction mechanisms. *Physiol. Meas.* 25:1077–1093.

99. Beebe, S. J., P. M. Fox, L. J. Rec, E. L. Willis, and K. H. Schoenbach. 2003. Nanosecond, high-intensity pulsed electric fields induce apoptosis in human cells. *FASEB J.* 17:1493–1495.
100. Nuccitelli, R., K. Tran, S. Sheikh, B. Athos, M. Kreis, and P. Nuccitelli. 2010. Optimized nanosecond pulsed electric field therapy can cause murine malignant melanomas to self-destruct with a single treatment. *Int. J. Cancer* 127:1727–1736.
101. Thomas, M. B., and A. X. Zhu. 2005. Hepatocellular carcinoma: the need for progress. *J. Clin. Oncol.* 23:2892–2899.
102. Costa, F., A. C. de Oliveira, R. Meirelles, T. Zanesco, R. Surjan, M. Chammas, et al. 2007. A phase II study of amplitude-modulated electromagnetic fields in the treatment of advanced hepatocellular carcinoma (HCC). *J. Clin. Oncol.* meeting abstract (25):15155.
103. Kirson, E. D., Z. Gurvich, R. Schneiderman, E. Dekel, A. Itzhaki, Y. Wasserman, et al. 2004. Disruption of cancer cell replication by alternating electric fields. *Cancer Res.* 64:3288–3295.
104. Nuccitelli, R., X. Chen, A. G. Pakhomov, W. H. Baldwin, S. Sheikh, J. L. Pomicter, et al. 2009. A new pulsed electric field therapy for melanoma disrupts the tumor's blood supply and causes complete remission without recurrence. *Int. J. Cancer* 125:438–445.
105. Majidian Eydgahi, S., J. Baharara, S. Zafar Balanezhad, and M. Asadi Samani. 2015. The synergic effect of glycyrrhizic acid and low frequency electromagnetic field on angiogenesis in chick chorioallantoic membrane. *Avicenna J. Phytomed.* 5:174–181.