

Membrane Patterns Carry Ontogenetic Information That Is Specified Independently of DNA

Jonathan Wells*

Biologic Institute, Redmond, Washington, USA

Abstract

Embryo development (ontogeny) depends on developmental gene regulatory networks (dGRNs), but dGRNs depend on pre-existing spatial anisotropies that are defined by early embryonic axes, and those axes are established long before the embryo's dGRNs are put in place. For example, the anterior-posterior axis in *Drosophila* and the animal-vegetal axis in *Xenopus* and echinoderms are initially derived from the architecture of the ovary through processes mediated by cytoskeletal and membrane patterns rather than dGRNs. This review focuses on plasma membrane patterns, which serve essential ontogenetic functions by providing targets and sources for intracellular signaling and transport, by regulating cell-cell interactions, and by generating endogenous electric fields that provide three-dimensional coordinate systems for embryo development. Membrane patterns are not specified by DNA sequences. Because of processes such as RNA splicing, RNA editing, protein splicing, alternative protein folding, and glycosylation, DNA sequences do not specify the final functional forms of most membrane components. Still less does DNA specify the spatial arrangements of those components. Yet their spatial arrangements carry essential ontogenetic information. The fact that membrane patterns carry ontogenetic information that is not specified by DNA poses a problem for any theory of evolution (such as Neo-Darwinism) that attributes the origin of evolutionary novelties to changes in a genetic program—whether at the level of DNA sequences or dGRNs. This review concludes by suggesting that relational biology and category theory might be a promising new approach to understanding how the ontogenetic information in membrane patterns could be specified and undergo the orchestrated changes needed for embryo development.

Cite as: Wells J (2014). Membrane patterns carry ontogenetic information that is specified independently of DNA. *BIO-Complexity* 2014 (2):1–28.
doi:10.5048/BIO-C.2014.2.

Editor: Ann Gauger

Received: January 27, 2014; **Accepted:** April 1, 2014; **Published:** May 22, 2014

Copyright: © 2014 Wells. This open-access article is published under the terms of the [Creative Commons Attribution License](#), which permits free distribution and reuse in derivative works provided the original author(s) and source are credited.

Notes: A *Critique* of this paper, when available, will be assigned **doi:**10.5048/BIO-C.2014.2.c.

*Email: jonwells2001@comcast.net

INTRODUCTION

In 1958—five years after he and James Watson discovered the molecular structure of DNA—Francis Crick proposed that the specificity of a DNA segment lies solely in its nucleotide sequence, which encodes the nucleotide sequence of a messenger RNA (mRNA) and thereby the amino acid sequence of a protein. Crick also proposed that the information encoded in DNA sequences can be transferred from DNA to protein, but not back again. He called the former the “sequence hypothesis” and the latter the “central dogma” of molecular biology [1], but many writers since that time have used “central dogma” to refer to the two proposals together. In this review I follow their practice.

Three years after Crick proposed the central dogma, François Jacob and Jacques Monod reported the discovery of a genetic switch in bacteria that regulates the synthesis of β -galactosidase, an enzyme (protein) needed to metabolize lactose. When the

bacteria are in a medium with no lactose, a DNA-binding protein represses the transcription of β -galactosidase mRNA, so the cells do not waste energy producing an unneeded protein. When lactose is present it induces the release of the repressor from the DNA, thus permitting the production of β -galactosidase. Jacob and Monod called this genetic system the “lac operon” [2,3].

In 1968 Eric Davidson published a book analyzing gene activity in early animal development that started with two premises: the central dogma (“Since the cell owes its definitive characteristics to the characteristics and functional attributes of its proteins... the differentiated state must ultimately depend on the transcription of genomic information”) and genomic equivalence (“Every living cell nucleus in a metazoan organism [with some exceptions, as Davidson noted] contains the same complete genome as was present in the zygote nucleus”).

Arguing from these two premises, Davidson concluded that animal embryogenesis (ontogeny) depends on “selective variation in gene activity” that involves regulatory mechanisms such as the lac operon [4].

Up until that time, many biologists had maintained that cytoplasmic factors are more important than nuclear genes in controlling ontogeny [5,6]. Indeed, in his 1968 book Davidson reviewed evidence that in several animal phyla “the initial, visible events of embryogenesis are *not* under the direct control of the embryonic cell genomes.” For example, the early development of cross-species hybrids in sea urchins as well as frogs follows the maternal pattern, with no detectable contribution from the paternal genome. Davidson concluded that early embryogenesis in these (and many other) animals is controlled by “morphogenetic determinants that are localized in the cytoplasm.” He emphasized, however, that the localized morphogenetic determinants are ultimately encoded by DNA; so are the transcription factors that regulate gene activity [4].

In 1970, Jacob wrote that an organism is the realization of a “programme génétique” (genetic program) written in DNA sequences [7]. The same year, Monod wrote that “the sequence of nucleotides in a DNA segment entirely defines the sequence of amino acids in the corresponding polypeptide [protein].” And since “the polypeptide sequence specifies completely (under normal initial conditions) the folded structure that the polypeptide adopts once it is constituted, the structural and hence functional ‘interpretation’ of genetic information is unequivocal, rigorous. No supplementary input of information other than the genetic is necessary; none, it seems, is even possible” [8]. Thereafter the idea of cytoplasmic control was largely replaced by the idea that genetic programs direct embryo development. A popular formulation of the central dogma is now “DNA makes RNA makes protein makes us” [9,10].

The emphasis on genetic programs owes much to evolutionary theory—specifically, to the modern synthesis of Darwinian evolution and Mendelian genetics. According to the modern synthesis, new heritable variations originate in genetic mutations. In a 1970 interview, Monod said that with the establishment of the central dogma, “and the understanding of the random physical basis of mutation that molecular biology has also provided, the mechanism of Darwinism is at last securely founded” [11].

Yet even in 1970 it was clear that there is no simple correspondence between DNA sequences (the genotype) and an organism’s observable characteristics (its phenotype), and therefore that the notion of “genetic program” needed—at the very least—further work. In what follows I summarize the subsequent elaboration of that notion in Eric Davidson’s idea of “gene regulatory networks.” I then survey evidence for the temporal and causal priority of spatial anisotropies in early development, and I argue that ontogeny in multicellular animals (metazoa) depends significantly on sources of spatial information in cells that are not reducible to DNA sequences. In particular, I focus on spatial information in plasma membranes, in the form of patterns of intracellular targets, complex carbohydrates, and ion channels. I conclude by discussing the

implications of membrane information for evolutionary theory and our understanding of ontogeny.

GENE REGULATORY NETWORKS

In 1969, Stuart Kauffman wrote that the analogy between digital computing and the control system exemplified by the lac operon had already suggested to several authors “that the genome embodies complex switching circuits which constitute a program for metabolic stability and cell differentiation.” Treating “genes” as binary switches, Kauffman used Boolean logic to model what he called “genetic nets” or “genetic regulatory circuits” [12].

In 1990, Davidson applied the term “gene regulatory networks” (GRNs) to sets of interacting DNA sequences, RNAs, and proteins that regulate transcription and other cellular processes. He argued that “the significant features” of early development in animal embryos “have to do directly with the distribution in embryonic space of gene regulatory molecules.” Davidson acknowledged that the distribution depended on spatial anisotropies in early embryos, but he considered those anisotropies less significant than their implementation by genes and gene products: “Since the basic objective is to understand what makes the various spatial domains of the early embryo functionally diverse, in terms of cell structure, behavior and synthetic activity, the problem of regulatory architecture has to be considered in particular at the level of the histospecific [i.e., tissue-specific] structural genes that endow the embryo with its specific regional functions” [13].

Since 1990 many authors have dealt with gene regulatory networks, but Davidson has been the most prolific writer on the topic. In 2006, he wrote that animal embryos illustrate two features:

The less important is the variable specifics of the initial cytoplasmic bases of spatial anisotropy. The other feature is of ultimate importance: This is the common functional endpoint of these very diverse initial stratagems for the spatial indication of future developmental domains. The principle is that whatever the bases of the anisotropies, however they come into being, whatever the cell fates that derive from what they set in train, they end up causing certain maternal transcription factors to be present and active in some spatially defined embryo nuclei, but not in others [14].

So Davidson maintained that GRNs are more important in ontogeny than the spatial anisotropies that precede them. In his view, the genetic programs embodied by GRNs control development, and mutations in the underlying genes lead to evolution. “Since the morphological features of an animal are the product of its developmental process,” he wrote, “and since the developmental process in each animal is encoded in its species-specific regulatory genome, then change in animal form during evolution is the consequence of change in genomic regulatory programs for development” [14].

Despite this emphasis on the genome, Davidson made it clear in 2011 that he was not advocating “the classic neo-Darwinian concept that evolution of animal morphology occurs by means of small continuous changes in primary protein sequence,” nor was he claiming that “evolution at all levels can be illuminated by detailed analysis of *cis*-regulatory changes in genes that are direct targets of sequence level selection.” These perspectives, he pointed out, “often focus on changes at single gene loci, and both are framed within the concepts of population genetics.” Instead, according to Davidson “the evolution of animal body plans is a system level property of developmental gene regulatory networks (dGRNs) which control ontogeny of the body plan” [15].

SPATIAL ANISOTROPIES IN ONTOGENY

Where do spatial regulatory domains come from? Far from being determined by the gene regulatory molecules that occupy them, as Davidson would argue, spatial domains determine the distribution of regulatory molecules. Spatial anisotropies precede—and are causally upstream of—the embryo’s dGRNs.

Drosophila

Among the best-studied dGRNs are those involved in the development of the fruit fly *Drosophila melanogaster*. The earliest dGRNs in ontogeny are customarily taken to consist of the “maternal effect” genes, which are involved in elaborating the anterior-posterior and dorsal-ventral axes of the oocyte [16–18]. For the anterior-posterior axis, maternal effect genes are involved in setting up a gradient of Bicoid protein with its highest concentration at the anterior end, and a gradient of Nanos protein with its highest concentration at the posterior end [19–21]. Other dGRNs are subsequently involved in establishing the dorsal-ventral axis, setting up body segments (“segmentation genes”), and specifying segment identities (“Hox genes”) [22–24].

The Bicoid gradient is commonly taken to be the starting-point of the first dGRN in *Drosophila* development [14,25]. The Bicoid gradient depends on a gradient of maternal *bicoid* mRNA that is highest at the anterior end of the oocyte [26,27]. When *bicoid* mRNA enters the oocyte from the nurse cells a microtubule-based mechanism concentrates it at the anterior end, where it becomes closely associated with the cell cortex

[28–30]. As used here, “cortex” refers to the plasma membrane and an underlying layer of cytoplasm consisting primarily of cytoskeletal elements (e.g., actin microfilaments and spectrins) [31]. The cortical localization of *bicoid* mRNA also requires several proteins, especially the endosomal sorting protein ESCRT-II [32,33].

Yet the anterior pole has to be in place before *bicoid* mRNA can be localized there. In fact, the *Drosophila* oocyte acquires its anterior-posterior polarity much earlier. The oocyte starts out in a follicle consisting of sixteen interconnected cells, fifteen of which will become “nurse cells” that synthesize and transport maternal mRNAs into the developing oocyte. Anterior-posterior polarity first appears in the oocyte when it becomes distinct from the nurse cells (Fig. 1) [34]. The selection of one cell to become the oocyte involves polarization of the membrane and depends on the asymmetric segregation of a germline-specific organelle called the “fusome,” an intracellular structure containing membrane-associated cytoskeletal components (such as spectrins) normally found in the cell cortex [35,36]. The fusome organizes the microtubule network early in oogenesis and directs the migration of centrosomes from the nurse cells into the new oocyte [37,38]. Intercellular signaling between the oocyte and the surrounding follicle cells then regulates the anterior-posterior patterning of the latter [39].

At first, the follicle containing the oocyte and its associated nurse cells is roughly spherical. Then molecular motors in a surrounding epithelial “corset” force the follicle to rotate several times, turning it into an oval with the oocyte at one end [40]. By this time, the anterior-posterior axis of the oocyte is well established—long before maternal effect gene products enter it from the nurse cells (Fig. 1, far right). So the anterior-posterior body axis in the *Drosophila* oocyte does not originate in maternal effect genes and their products, but in spatial asymmetries inherent in the architecture of the ovary and in polarized membranes and cytoskeletal components of the follicle [41].

Xenopus

In the frog *Xenopus laevis*, oocytes have neither anterior-posterior nor dorsal-ventral axes. Instead, a *Xenopus* oocyte has an animal-vegetal axis, with the animal hemisphere covered by dark pigment granules and the vegetal hemisphere filled with large yolk platelets. After fertilization the dorsal-ventral axis




Figure 1: *Drosophila* oogenesis. The *Drosophila* oocyte originates in a follicle consisting of sixteen interconnected cells. Polarization of the cytoskeleton and membrane specify one cell to become the oocyte; its fifteen neighbors then become nurse cells that synthesize maternal mRNAs. The follicle is roughly spherical at first (left); molecular motors in the surrounding epithelium then rotate it, forming an oval with the oocyte at one end. By this time the anterior-posterior axis of the oocyte is well established, before maternal mRNAs are transported into it from the nurse cells (right).

doi:10.5048/BIO-C.2014.2.f1

is established by a process of “cortical rotation,” in which the cortex rotates approximately 30° relative to the interior of the zygote [42,43]. The movement is microtubule-dependent, and the cortex normally rotates away from the sperm entry point, which has suggested to some that the aster nucleated by the sperm centrosome initiates cortical rotation [44]. The rotation movement orients microtubules in the vegetal cortex, and these form a parallel array that provides tracks to transport various proteins from the vegetal pole to the future dorsal side of the embryo, thereby establishing a dorsal-ventral axis [45,46]. One of the most important of these proteins is β -catenin, a member of the Wnt signaling pathway that is involved in cell-cell communication and gene regulation in the early development of many animals, including *Drosophila* and *Xenopus* [47,48]. The cortical region in which these proteins accumulate subsequently becomes the “blastopore,” an opening through which cells migrate inward during gastrulation to establish the anterior-posterior axis. The dorsal lip of the blastopore has been called the “organizer,” because transplanting it to the presumptive future ventral side of another embryo induces the latter to organize a second tadpole conjoined to the first [49].

The dorsal-ventral and anterior-posterior axes of the *Xenopus* embryo depend on the prior establishment of the animal-vegetal axis in the oocyte. *Xenopus* oogenesis is customarily divided into six stages [50]. From stage II through stage V, maternally produced yolk is transported across the entire surface of the developing oocyte. Once inside, yolk platelets are concentrated in the vegetal hemisphere by intracellular transport (Fig. 2) [51]. More than twenty RNAs are also concentrated in the vegetal hemisphere, by two different mechanisms exemplified by the maternal RNAs *Vg1* and *Xcat-2* [52]. *Vg1* is subsequently involved in mesoderm induction during embryogenesis, and *Xcat-2* (which has some sequence similarity to *Drosophila nanos*) may be subsequently involved in germ cell development [53,54]. During stage IV, *Vg1* is transported to the vegetal pole by a microtubule-based mechanism and then anchored to the vegetal cortex by a mechanism requiring membrane-associated actin microfilaments [55]. But earlier in oogenesis a microtubule-independent mechanism localizes *Xcat-2* RNA to the vegetal pole along with a “mitochondrial cloud,” also known as the “Balbiani body” [56,57].

The Balbiani body is an aggregate of mitochondria, cytoskeletal elements, endoplasmic reticulum, RNAs and

proteins [58]. In early oocytes it is located next to the germinal vesicle (the maternal pronucleus) and is the first indicator of animal-vegetal polarity [59,60]. In this respect it is like the *Drosophila* fusome; indeed, a fusome entering a *Drosophila* oocyte resembles a Balbiani body [61]. During *Xenopus* oogenesis, RNAs diffuse into the Balbiani body and are trapped there by elements of the endoplasmic reticulum [62]. The Balbiani body then expands directionally toward the vegetal pole by an unknown mechanism, delivering RNAs such as *Xcat-2* to the vegetal cortex [63].

In the microtubule-dependent transport that occurs later (in stage IV), the localization of *Vg1* RNA to the vegetal pole depends on a “localization signal” in the RNA’s 3’ untranslated region [64,65]. In this respect, the localization of *Vg1* is similar to localization of other mRNAs in a variety of organisms, including *bicoid* in *Drosophila* [66]. The *Vg1* localization signal works, however, only because its cortical target is already in place. Once RNAs are anchored at the vegetal pole of a *Xenopus* oocyte, isolated pieces of the vegetal cortex retain them [67]. The same is true of the dorsal cortex of *Xenopus* zygotes after cortical rotation [68]. Some RNAs might be anchored to cytoskeletal elements associated with the cortex (such as actin microfilaments), but evidence indicates that at least some RNAs are anchored to the vegetal cortex in a membrane-dependent manner [69].

Echinoderms

The oocytes of echinoderms (e.g., sea urchins and starfish), like those of frogs, have a primary animal-vegetal axis [70,71]. The second axis to form is the oral-aboral axis, which extends from the mouth to the opposite side of the organism [72]. Exactly when the oral-aboral axis forms and how it is determined have been controversial and vary from species to species [73–81].

Since the adults of most echinoderms exhibit five-fold radial symmetry, the identification of anterior-posterior and dorsal-ventral body axes in these animals has been problematic. The oral-aboral axis is commonly equated to the dorsal-ventral axis, because it is the second to form and most species orient with the mouth down, on the ventral side [74,78,79,82]. But the oral-aboral axis has also been equated to the anterior-posterior axis, because in bilaterally symmetrical animals the mouth is at the anterior pole and echinoderms are thought to have evolved from bilaterian ancestors [83,84].




Figure 2: *Xenopus* oogenesis. An early *Xenopus* oocyte (left) is approximately spherically symmetrical. As yolk is transported into it from the surrounding ovary, it becomes more polarized. The different shades of gray in this simplified drawing represent layers of yolk; the lighter areas are younger. The clear area is the germinal vesicle (the maternal nucleus), which becomes more elliptical as the oocyte matures. Animal-vegetal polarity is clearly visible in the distribution of the yolk and the location of the germinal vesicle near the animal pole (second from left), before maternal mRNAs enter the oocyte (right) to be differentially distributed along the animal-vegetal axis. doi:10.5048/BIO-C.2014.2.f2

In any case, it is clear that an echinoderm oocyte has an animal-vegetal axis that precedes the formation of the oral-aboral axis and is necessary for proper development. When unfertilized *Paracentrotus lividus* or *Hemicentrotus pulcherrimus* eggs are surgically bisected in a plane that includes the animal-vegetal axis, both halves proceed through gastrulation, but when the eggs are bisected in a plane perpendicular to the axis only the vegetal half undergoes development past the blastula stage [73,85]. Since β -catenin is involved in vegetal development, it has been suggested that it accumulates at the vegetal pole of the oocyte while other molecules localize to the animal pole [86–89]. Several specific mRNAs are also differentially localized in the oocyte with respect to the animal-vegetal axis [90,91].

Just as in *Drosophila* and *Xenopus* oocytes, the localization of specific molecules to the animal and vegetal poles in echinoderm oocytes presupposes that those poles are already present. In several sea urchin species, the animal pole of a primary oocyte is marked by a “jelly canal” that admits foreign ink particles excluded elsewhere from the cell [70,85,92,93]. In immature starfish oocytes, the animal poles are distinguished by the adjacent eccentric germinal vesicle (the maternal nucleus) and centrosomes of the pre-meiotic aster [94,95]. In any case, it is clear that the animal-vegetal axis of an echinoderm oocyte is present at the earliest stage of oogenesis and presumably derived from the architecture of the ovary.

Furthermore, the animal-vegetal axis is clearly expressed in the cortex. In starfish, the patterning of the cortex depends on microtubules arrays [95,96]. In sea urchins, the oocyte cortex contains arrays of microfilaments, microtubules, and intermediate filaments [97,98]. So in sea urchins and starfish, as in *Xenopus laevis* and *Drosophila melanogaster*, some spatial information in oocytes is imparted by polarized cytoskeletal arrays, and some is imparted by patterns in the plasma membrane or cortex.

The cytoskeleton and plasma membrane interact in complex ways, so the patterns in one cannot be completely understood apart from patterns in the other. Nevertheless, the remainder of this review focuses primarily on patterns in the plasma membrane.

BIOLOGICAL MEMBRANES

In 1925, Evert Gorter and François Grendel proposed that biological membranes are composed of lipid bilayers, with the hydrophobic ends of the constituent lipid molecules embedded in the middle of the bilayer [99]. In 1970, L. D. Frye and Michael Edidin labeled cell surface antigens in mouse cells with an antibody that fluoresced at a particular wavelength, and they labeled cell surface antigens in human cells with an antibody that fluoresced at a different wavelength. Then they used a virus to induce fusion of the mouse and human cells into heterokaryons. After about 40 minutes at 37° C, the labeled antigens had intermixed, spreading over the entire membrane in most of the heterokaryons. When Frye and Edidin inhibited protein synthesis there was no effect on the rate of intermixing, but when they lowered the temperature to 15° C the rate decreased. The results were consistent with simple diffusion, and

Frye and Edidin concluded that membrane-associated proteins are free to diffuse laterally in the plane of the membrane [100].

Citing the work of Frye and Edidin two years later, Seymour Singer and Garth Nicolson proposed a “fluid mosaic model” of cell membranes. In their model, the lipid bilayer is like a two-dimensional sea in which mostly monomeric proteins float, unrestricted in their lateral movements except for isolated local interactions [101]. But a new technique called fluorescence recovery after photobleaching (FRAP) soon provided evidence that the fluid mosaic model was inadequate. In the FRAP technique, a membrane is first labeled with fluorescent probes; a highly focused bright light bleaches the probes in a small patch of the membrane; and fluorescent molecules then diffuse into the bleached patch [102,103]. Results showed that the rate of diffusion of many proteins in natural (as opposed to artificial) biological membranes was much slower than predicted by the fluid mosaic model [104,105].

In 1984 Wolfhard Almers and Charles Stirling reported, “It is becoming clear that in fully differentiated tissues, a large fraction of membrane protein is neither freely mobile nor randomly distributed... Evidently, Frye and Edidin’s 1970 finding of unrestricted mobility was atypical” [106]. Evidence against unrestricted mobility continued to accumulate, and in 1991 Edidin (along with Scot Kuo and Michael Sheetz) concluded that “the lateral diffusion of most [membrane] proteins is hindered in several ways,” producing spatially restricted domains or “patches” rather than a random distribution [107].

In 2005, Donald Engleman concluded that the fluid mosaic model needed to be re-examined, because each of its principal ideas was “misleading.” Instead of being a lipid bilayer sea in which widely dispersed monomeric proteins float freely, biological membranes are “typically crowded” with oligomers and “preferential associations,” such that “patchiness is the order of the day” [108]. The same year, Akihiro Kusumi and his colleagues modeled the plasma membrane “as a compartmentalized fluid, in which compartmentalization is caused by the fence (corralling) effects of the membrane skeleton [which is actin-based in most mammalian cells] as well as the hydrodynamic slowing effects of transmembrane-protein pickets anchored on the membrane-skeleton fence” [109].

Proteins are not the only molecules heterogeneously distributed in the membrane—lipids are, too. Various researchers in the 1970s and 1980s presented evidence for spatial domains in membrane lipids [110–112]. Domains rich in sphingolipids and cholesterol were called “lipid rafts” [113]. For several years their existence was controversial [114–117], but improved experimental techniques subsequently yielded abundant evidence for lipid raft nanodomains [118–122]. Like membrane protein compartments, lipid rafts may be formed and maintained, in part, through cytoskeletal interactions [123,124], and they play important roles in membrane organization and signal transduction [125–127].

So biological membranes are patterned in complex ways. Those patterns serve important functions in cells, tissues and embryos. The following sections summarize the roles of plasma membrane patterns in (a) providing targets and sources for intracellular transport and signaling, (b) regulating cell-cell

interactions by means of a “sugar code,” and (c) generating endogenous electric fields that provide three-dimensional coordinate systems for ontogeny.

INTRACELLULAR TARGETS AND SIGNALING

The precise intracellular localization of mRNAs is functionally important in many animal cells—not just oocytes [128–131]. The localization of mRNAs commonly depends on specific sequences in their untranslated regions that have been called “zipcodes” [132]. Like postal zipcodes, such sequences identify the “addresses” in the cell to which the mRNAs are to be sent. Like a postal zipcode, however, an mRNA zipcode is meaningless unless it matches a pre-existing address—that is, a target.

Evidence from a variety of cells suggests that mRNA localization requires the binding of a protein to the zipcode to form a ribonucleoprotein particle (RNP); the combination is then transported to its destination [133]. A well-studied example is the RNA zipcode-binding protein ZBP-1, which is necessary for the localization of β -actin mRNA to the leading edges of migrating fibroblasts (where actin microfilaments are polymerizing) [134–136]. Another example is Vg1 RBP (for “RNA-binding protein”), which is necessary for localizing *Vg1* mRNA to the vegetal pole in *Xenopus* oocytes [65,137]. Still another example is heterogeneous nuclear ribonucleoprotein A2 (hn-RNP A2), which is involved in localizing mRNAs in neurons [138].

Like zipcodes themselves, however, zipcode-binding proteins do not specify the destination. Using the postal code metaphor, zipcode-binding proteins could be likened to cargo containers, cytoskeletal motor molecules to delivery trucks, and the cytoskeleton to the highway system on which the trucks travel. But destinations for intracellular transport—like the geographical addresses in a postal delivery system—must also be specified.

In some cases, destinations might be specified by the spatial arrangement of microtubules; in the postal metaphor, packages could be dispatched on a particular highway and then carried to the end of the road and simply dropped off. In some cases, however, destinations are known to be specified by targets in the form of membrane-bound proteins that respond to extracellular cues. For example, β -actin mRNA localization in fibroblasts is induced by receptor tyrosine kinases (RTKs) in the plasma membrane that respond to extracellular platelet-derived growth factor [139,140]. RTKs consist of a ligand-binding extracellular domain, a single α -helix that passes through the membrane, and an intracellular domain containing a protein tyrosine kinase [141,142]. Upon activation, the tyrosine kinase transfers a phosphate group from ATP to a protein, initiating a complex and dynamic intracellular signaling cascade that includes small guanosine-triphosphatases (GTPases) of the Rho family. These molecules—especially the cell division control protein Cdc42—polarize the microtubule network that delivers β -actin mRNA to the source of the signaling cascade [143–146]. The fibroblast then migrates in the direction of the external cue.




Figure 3: Comparison of mesenchymal and epithelial polarity. The illustration A) shows a highly simplified drawing of a mesenchymal cell moving along extracellular matrix (the heavy straight line at the bottom). Activated membrane-bound receptor tyrosine kinases (green) induce front-to-back polarity and actin polymerization at the leading edge. The illustration B) shows a highly simplified epithelial cell, resting on the extracellular matrix and tightly bordered on either side by other epithelial cells (not shown). Channels in the basal membrane (red) pump sodium ions out of the cell and channels in the apical membrane (blue) admit sodium ions into the cell. The result is a trans-epithelial potential, basal side positive. In both cells the large circle in the center is the nucleus, the parallel stack of inner membranes is the Golgi apparatus, and the straight lines are microtubules. doi:10.5048/BIO-C.2014.2.f3

In an animal embryo, cells that migrate easily (such as fibroblasts) are called mesenchyme, while cells that are immobilized in adherent sheets are called epithelium. Mesenchymal cells exhibit front-to-back polarity, with an axis oriented in the direction of movement. Immobilized epithelial cells exhibit top-to-bottom (apical-basal) polarity, with an apex oriented toward a lumen or the outside of a tissue and a base attached to an extracellular substratum (Fig. 3). Cells in epithelia are also attached to each other laterally by tight junctions to form an impervious protective layer [147,148].

Mesenchymal cells can transform into epithelial cells, and *vice versa*, in what are called the mesenchymal-epithelial transition and the epithelial-mesenchymal transition, respectively [149]. Both transitions are essential in metazoan ontogeny. Organizing cells into tissues depends on coordinating cell polarities, and in the process cells can switch between various polarized states [150]. After gastrulation, vertebrate embryos develop a neural tube composed of epithelial cells. At the end of neurulation, some of these (the neural crest cells) transform into mesenchymal cells that migrate elsewhere in the embryo to form craniofacial structures, the peripheral nervous system, pigment cells in the skin, and parts of the cardiovascular system [151,152]. Upon reaching their destinations, most of these mesenchymal cells then transform back into epithelial cells [153].

Polarity in an epithelial cell is normally initiated by contact with another cell [154]. The contact point produces an asymmetry in the plasma membrane that leads to the asymmetric localization of various proteins [155]. This asymmetric localization is mediated by differences in phosphorylated lipids in the apical and basal membranes [156,157], and by various sorting signals associated with the proteins. A particularly well-studied example of the latter is the glycoposphatidylinositol “anchor” [158,159]. Two key signaling molecules are the phosphoinositides $\text{PtdIns}(3,4,5)\text{P}_3$ and $\text{PtdIns}(3,4)\text{P}_2$. In mesenchymal cells, these regulate protrusion of the front




Figure 4: Linear vs. branched information. Illustrated are A) a trinucleotide, B) a tripeptide, and C) a trisaccharide. Nucleotides in DNA are covalently linked to each other only in linear chains; the same is true for amino acids in polypeptides (proteins), with a few exceptions. But monosaccharides can be covalently linked to each other through one or more of their hydroxyl groups, so they can be combined with other monosaccharides in several different ways to form branched glycans. [doi:10.5048/BIO-C.2014.2.f4](https://doi.org/10.5048/BIO-C.2014.2.f4)

and retraction of the back to produce forward movement. In epithelial cells, the same molecules maintain the functional differences between the apical and basal domains [160,161].

Although polarity in mesenchymal and epithelial cells is normally initiated in response to extracellular cues, the plasma membranes of many other animal cells have spatially heterogeneous patterns that are more intrinsic to isolated cells.

THE SUGAR CODE

The plasma membranes of all living cells studied to date are covered by arrays of carbohydrates called “glycans” [162]. Glycans can be attached to lipid molecules (glycolipids) or to proteins (glycoproteins), and many of them are quite complex [163]. In living cells, nucleotides in DNA are covalently linked to each other in linear chains; with some exceptions, the same is true for amino acids in proteins. But monosaccharides can be covalently linked to each other through one or more of their hydroxyl groups. Since D-glucose (for example) has five hydroxyl groups, one of which can assume two different positions, it can be covalently linked to other monosaccharides in six different ways (Fig. 4). As a result, carbohydrates can form branching chains that are far more elaborate than linear chains

of nucleotides and amino acids (Fig. 5) [164].

While the four nucleotides in the genome can form a maximum of $4^6 \approx 4 \times 10^3$ hexanucleotides, and the twenty amino acids in the proteome can form a maximum of $20^6 \approx 6 \times 10^7$ hexapeptides, the dozen or so monosaccharides in the “glycome” can theoretically form more than 10^{12} hexasaccharides. Clearly, the information-carrying capacity of the “glycome” far exceeds the combined capacities of the genome and the proteome. The information carried by the glycome has been called the “glycocode” or “sugar code” [165–169].

Glycosaminoglycans (GAGs) are unbranched polysaccharides composed of disaccharide subunits containing an amino group. Yet although they are unbranched, they can be assembled from dozens of different subunits, and sulfate groups can be attached to them in a wide variety of patterns. For example, a sulfate group can be attached to a trisaccharide in ten different positions, increasing its information-carrying capacity tenfold [170]. This makes GAGs some of the most information-dense molecules in biology [171–173]. Of the five types of glycosaminoglycans, four are covalently attached to proteins to form proteoglycans (PGs). Like glycolipids and glycoproteins, PGs are common in the plasma membranes of many cells.

The complexity of glycans has made it difficult to study them in living cells and tissues with standard biochemical methods, but antibodies have proven useful in identifying specific carbohydrate moieties [174]. Studies using monoclonal antibodies have shown that cell-surface glycans in early mouse embryos change in a highly ordered and stage-specific manner; the data suggest that they mediate cellular orientation, migration, and responses to regulatory factors during development [175–177]. Monoclonal antibody studies have also found cell- and stage-specific changes in cell-surface glycans during early embryogenesis in the chick [178,179], and during neural development in late-stage rat embryos [180].

Genetic analyses have produced comparable results. Mutations in genes encoding the protein portions of glycoproteins and sulfated PGs, and mutations in genes encoding various enzymes involved in glycosylation or sulfation, have revealed spatiotemporal expression patterns in the embryos of *Caenorhabditis elegans*, *Drosophila melanogaster*, mice and humans [181,182], as well as in zebrafish [183]. These patterns play important roles in development [184–186].




Figure 5: A branched glycan. The five different shapes here represent five different monosaccharides. This shows one way these monosaccharides could be covalently connected to form a branched glycan. The “X” on the left could be either a protein (in which case this would be a glycoprotein) or a lipid molecule (in which case this would be a glycolipid). More monosaccharides (including others not represented here) could be attached on the right (among other places), yielding trillions of different glycans carrying different glycocodes. [doi:10.5048/BIO-C.2014.2.f5](https://doi.org/10.5048/BIO-C.2014.2.f5)

Another approach to studying cell-surface glycans is matrix-assisted laser desorption/ionization (MALDI) mass spectrometry imaging [187]. This technique has been used to study gangliosides, a class of glycolipids that are typically partitioned into lipid rafts and have hundreds of different carbohydrate side-chains. MALDI mass spectrometry has revealed striking differential distributions of gangliosides in mouse brain tissues [188].

The sugar code can be “interpreted” by proteins called lectins. Unlike antibodies, lectins are not produced by the immune system, and unlike enzymes they do not catalyze biochemical reactions, but like antibodies and enzymes they “recognize” specific three-dimensional structures of other molecules. They do this by means of “carbohydrate recognition domains” (CRDs) [189–191].

Lectins mediate a wide variety of cell-cell interactions [192–194]. They have also been useful in studying the distribution and functions of cell-surface glycans. For example, lectins with different specificities have revealed spatial and temporal changes in the distribution of cell surface oligosaccharides during gastrulation and neurulation in chick embryos [195]. Comparative analyses with lectins have also shown reproducible differences in the spatial distribution of cell-surface carbohydrates in the olfactory sensory neurons of mice [196].

Galectins are a taxonomically widespread family of animal lectins that bind glycans containing β -galactoside [197–199]. Certain galectins are differentially expressed in early human embryogenesis, suggesting a role in tissue differentiation [200]. Other galectins are temporally and spatially regulated in frog embryos [201]. By forming lattices with glycoproteins on the plasma membrane, galectins help to organize cell-surface domains and thereby affect cell signaling [202,203]. Recent work has shown that a network of two galectins interacting with cell-surface moieties regulates skeletal morphogenesis in chick limbs [204].

The examples cited above involve tissue-level differences, but spatial localization of cell-surface glycans has also been observed in single cells. Several examples involve fungal cells: Immunological analyses have demonstrated changes in cell wall glycoprotein distribution during germ-tube formation in *Candida albicans* [205]; *Cryptococcus neoformans* cells exposed

to fluorescently labeled lectin showed marked localization of cell-surface glycans around sites of cell division [206]; and a similar approach revealed heterogeneous subpopulations of lectin ligands on hyphal bodies of *Beauveria bassiana* [207].

Studies using fluorescently labeled lectins have shown that several glycosidases—enzymes that catalyze the hydrolysis of the bonds joining carbohydrates to other molecules—occur in the plasma membrane of *Drosophila* spermatozoa. The glycosidases are localized to the sperm’s acrosomal region (the tip of the head) and the tail [208,209]. Complementary carbohydrate residues have been identified in mature *Drosophila* oocytes, where they are localized to the micropyle—a hollow anterior projection through which the sperm enters. This pattern of cell surface glycans suggests a role for them in fertilization (Fig. 6) [210–212].

New techniques may soon make it possible to characterize more precisely the spatial localization of specific glycans on the surfaces of single cells. For example, labeled sugars can be incorporated into specific carbohydrates in a process called “metabolic oligosaccharide engineering” [213–215]. This technique has been used to study spatiotemporal differences in cell-surface glycans during zebrafish development [216]. Another potentially fruitful approach is to use functionalized quantum dots and glyconanoparticles for the fluorescent imaging of dynamic glycan expression on single cells [217,218].

ENDOGENOUS ELECTRIC FIELDS

It has long been known that probably all living cells (not just nerve and muscle cells) generate electric fields across their membranes [219–222]. In animal cells, a sodium-potassium pump in the membrane utilizes energy from ATP to move three sodium ions out of the cell while taking in two potassium ions [223,224]. This raises the intracellular concentration of potassium ions, which corrects the imbalance by flowing out of the cell through ion-selective channels in the membrane. The combined action of sodium-potassium pumps and potassium “leak” channels makes the interior of the cell electrically negative with respect to the exterior. The resulting voltage difference across the membrane is called the “membrane potential” [225–227].

The voltage difference (V) per unit distance is the electric



Figure 6. Glycans in *Drosophila* fertilization. This shows a *Drosophila melanogaster* spermatozoon and oocyte just before fertilization. The two are drawn to scale: The sperm is ~ 1.8 mm long, and the oocyte is ~ 0.5 mm long. The acrosome (the tip of the head of the sperm) contains the glycosylated enzyme α -L-fucosidase (green). The portion of the sperm head just behind the green label contains the nucleus; farther back, the sperm tail contains more α -L-fucosidase (not shown). The micropyle (a small hollow projection at the anterior end of the oocyte through which the sperm will enter the egg) contains α -L-fucose (orange). The two complementary glycans are thought to mediate fertilization. doi:10.5048/BIO-C.2014.2.f6




Figure 7. Endogenous electric fields in *Xenopus* embryos at progressive stages of development. A *Xenopus* oocyte (A) is shown with some of the field lines of the endogenous electric field (EEF) surrounding it. Positive ions exit through the vegetal pole and enter through the animal pole. A *Xenopus* neurula (B) is about the same size as the oocyte but consists of thousands of cells. The field lines show some of the paths of positive ions around the neural fold, which will close to form the embryonic spinal column. A *Xenopus* tadpole (C) has positive ions leaving the future site of hindlimb development and entering through the gills, creating an endogenous electric field. doi:10.5048/BIO-C.2014.2.f7

field (E), which in biological systems is usually expressed in units of mV/mm. The direction of an electric field is taken to be the direction in which positive charges flow; in living cells and embryos, electric currents consist of positive and negative ions rather than electrons. The amount of electric current (A) flowing across a unit area is the current density (J), which in biological systems is usually expressed in units of $\mu\text{A}/\text{cm}^2$. Electric field and current density are quantitatively related by the equation $E = J\rho$, in which ρ is the resistivity of the medium, measured in ohm-centimeters ($\Omega\cdot\text{cm}$). Endogenous electric current densities are usually of the order of 1–100 $\mu\text{A}/\text{cm}^2$ [228].

Cellular membrane potentials generally range from ~ 20 to ~ 200 mV DC. Multicellular organisms, and their organs, are covered by epithelia, which (as we saw above) are polarized. Among other things, the ion channels in the apical domain are different from those in the basolateral domain [161]. The result is a “transepithelial potential” (TEP) that depends on sodium ions. Unlike the transmembrane potential of individual cells, the TEP is usually negative on the outside of the organ or organism and positive on the inside. Most TEPs range from 15 to 60 mV (though they can be lower or much higher), and endogenous electric fields (EEFs) are typically in the 10–150 mV/mm range [229–231].

Membrane potentials were first measured with sharp microelectrodes developed in the 1940s [232]. In this technique, a very small microelectrode is inserted into the cytoplasm of a cell and the potential difference is measured between it and a reference electrode in the surrounding medium. In the 1970s, Erwin Neher and Bert Sakmann developed the technique of patch clamping, which uses a small glass pipette in contact with (but not penetrating) the plasma membrane to measure the membrane potential [233]. Since a sharp microelectrode unavoidably causes some leakage through the membrane, patch clamping is preferred [234].

Transmembrane voltage can also be measured using voltage-reporting dyes. In Förster resonance energy transfer (FRET), a dye molecule (the “donor”) is stimulated to fluoresce at its characteristic wavelength; at the same time it transfers energy to a different dye molecule (the “acceptor”) adjacent to it, which fluoresces at a different wavelength. If the two molecules are separated by even a small distance, the energy transfer does not

occur and only the first molecule fluoresces. A useful donor is the phospholipid CC2-DMPE, which binds to the outside of the cell membrane, while a negatively charged hydrophobic acceptor such as DiBAC₄(3) can move across the cell membrane in response to changes in membrane potential. When the donor is stimulated, the ratio between the amounts of fluorescence at the dyes’ two wavelengths measures the degree of their separation and thus the membrane potential [235–237].

Endogenous direct current electric fields around cells and embryos can be measured with an ultrasensitive vibrating probe invented in 1974 by Lionel Jaffe and Richard Nuccitelli. The original probe measured the field difference between two points 30 μm apart while oscillating at 200 Hz to minimize noise [238]. Vibrating probes—enhanced by several technological improvements—are still being used to measure EEFs and ion currents in and around biological systems [239,240].

EEFs in oocytes and embryos

Mature *Xenopus* oocytes that have been dissected from their follicles have membrane potentials between 60 and 80 mV [241]. The resulting EEF is polarized along the animal-vegetal axis, and the oocytes drive electric currents with a density of about 1 $\mu\text{A}/\text{cm}^2$ through themselves, with positive charges entering around the animal pole and exiting through the vegetal pole (Fig. 7A) [242]. After fertilization the embryo enters the cleavage stage; without increasing its overall size, it subdivides into thousands of cells called blastomeres. In the first and second cleavage divisions, large electrical currents leave the new membrane forming in the furrows and return through the older membrane outside the furrow [243]. As a *Xenopus* embryo continues to cleave, it forms a fluid-filled central cavity called a blastocoele, and the blastomeres develop into an epithelium with an apical side facing the external medium. The resulting transepithelial potential is about 20 mV, blastocoele positive [244].

When cleavage ends gastrulation begins, and cells from the exterior migrate into the blastocoele through the blastopore (the location of which was previously determined by cortical rotation). A large current flows outward through the blastopore, carried mainly by Na^+ ions [245]. When the neural folds form, a steady ionic current is driven out of the lateral walls of the folds (Fig. 7B) [246]. After the folds close, the neural tube

maintains a potential difference of ~ 18 mV across its walls; since the walls are only ~ 40 μm thick, the resulting electric field across the neural tube cells is over 400 mV/mm [247]. The neurula then elongates into a tadpole, and electric currents flow outward from the two flank areas where the hind limbs will form and flow back inward through the gills (Fig. 7C) [248]. These currents reach their maxima (about 2 $\mu\text{A}/\text{cm}^2$) just after the limb buds become visible and decline after that [249].

Ionic currents are also present in developing chick embryos. The embryos of birds form a double-layered “blastodisc” of epithelial cells resting on the egg yolk; a groove called the “primitive streak” then forms in the outer layer. During gastrulation, cells migrate through the primitive streak into the space between the two layers. Measurements with a vibrating probe show that strong electrical currents leave the primitive streak, with densities of 10–20 $\mu\text{A}/\text{cm}^2$ [250]. The Na^+ -dependent currents return through the periphery of the blastodisc [251]. Even larger currents have been detected at the posterior intestinal portal (PIP), located at the tail end of the primitive streak. In chick embryos between two and four days old, currents with densities of 46–60 $\mu\text{A}/\text{cm}^2$ begin to flow into the PIP, followed later by outward currents with densities as high as 192 $\mu\text{A}/\text{cm}^2$ [252].

Early mammalian embryos also develop as blastodiscs and form primitive streaks. Ionic currents with densities of 20–60 $\mu\text{A}/\text{cm}^2$ have been detected leaving the primitive streaks of mouse embryos [253]. Like frog embryos, mouse embryos produce outward currents associated with their limb buds. Chick embryos also produce very large outward currents (about 100 $\mu\text{A}/\text{cm}^2$) at the future sites of their limb buds, three developmental stages before the latter become visible [254].

Effects of electric fields in vitro

How might embryonic electric currents and fields influence development? One way might be to direct cell movements. For over a century electrically guided locomotion, called “galvanotaxis,” has been observed in cells from a variety of organisms in the presence of artificially applied electric fields [255]. But can relatively small *endogenous* electric fields produce galvanotaxis? One way to test this is to apply artificial electric fields of physiological strength to embryonic cells *in vitro*.

In the 1980s, Robert Stump and Kenneth Robinson reported that *Xenopus* neural crest cells migrated toward the cathode (the negative pole) of an applied DC electric field of 10 mV/mm, well within the range of observed EEFs [256,257]. Richard Nuccitelli and Carol Erickson reported the same behavior for quail neural crest-derived fibroblasts [258,259]. By contrast, chick embryonic Schwann cells (which are derived from peripheral nerves) migrate preferentially toward the anode (the positive pole) in DC electric fields ranging from 10 to 100 mV/mm [260]. Not all embryonic cells, however, migrate directionally in an electric field of physiological strength [261,262].

Applied electric fields can also affect neural networks. Nerve cells establish contact with each other by extending projections called neurites. When embryonic chick ganglia were placed

in DC electric fields ranging from 70–140 mV/mm, their neurites grew faster toward the cathode than the anode [263]. Embryonic *Xenopus* neurons reportedly do the same in fields as low as 7 mV/mm [264,265].

Disrupting bioelectric patterns in embryogenesis

The most compelling evidence that ion currents, transmembrane voltage potentials and EEFs play significant roles in ontogeny comes from artificially disrupting them *in vivo* and then observing the effects of their disruption on morphogenesis. As we saw above, a strong current flows outward from the posterior intestinal portal (PIP) in four day-old chick embryos. In 1992, Kenneth Robinson and Kevin Hotary implanted conductive shunts into such embryos and thereby reduced by 30% the endogenous electric currents leaving the PIP. Over 90% of the affected embryos exhibited developmental abnormalities, most (as expected) in tail development—ranging from abnormal morphology to complete absence. In a control group receiving non-conductive implants, only 11% displayed defects [266].

In gastrulating *Xenopus* embryos, as we saw above, a large current flows outward through the blastopore. This current is associated with an intraembryonic head-to-tail EEF of 27 ± 4 mV/mm. In 1994, Hotary and Robinson inserted microelectrodes into embryos at this stage and applied electric currents of various magnitudes. A group of twenty-three embryos received currents of 100–500 nA in the opposite direction of the normal outward current; of these, twenty developed abnormalities that included reduced heads and retarded eye formation. Microelectrodes were also inserted in a control group of fourteen embryos, but the applied current was either 10 nA or zero; of these, only one developed abnormally [267].

After gastrulation, as we saw above, the neural tube of a *Xenopus* embryo maintains a potential difference across itself. In the mid-1990s, Riya Shi and Richard Borgens studied the transneural tube potential (TNTP) in axolotl embryos and reported it to be 40–90 mV. As in *Xenopus*, the lumen of the neural tube in axolotls is negative with respect to the outside. When Shi and Borgens injected Na^+ channel blockers into the lumen through microelectrodes, the TNTP dropped by 30% immediately and by 80% within fifteen minutes. Of fifty embryos treated in several different experiments all but one developed severe cranial defects, ranging from substantial deletions of central nervous system structures to the complete absence of a head. In one control group, eleven embryos were similarly injected, but without Na^+ channel blockers; all developed normally. As another control, the Na^+ channel blockers were injected just under the surface ectoderm (rather than into the neural tube) in twenty embryos, and all but one of these developed normally [268,269].

In 2011 Laura Vandenberg, Ryan Morris and Dany Adams disrupted cranial development in *Xenopus* tadpoles by inhibiting an ion channel protein. Different subpopulations of ectodermal cells in normal tadpoles are distinguished by different membrane voltages generated by ATP-dependent proton pumps in the

plasma membrane known as (H⁺)-ATPases (or V-ATPases, because they were originally discovered in vacuolar membranes) [270–272]. Adams and her colleagues had previously shown that V-ATPases are necessary in left-right patterning and tail regeneration in *Xenopus* [273,274]. V-ATPases are also involved in the anterior-posterior patterning of the *Xenopus* nervous system [275]. Vandenberg and her colleagues made a mutant mRNA for ductin, the proton-pumping subunit of V-ATPases [276]. This mRNA was translated into non-proton-pumping ductin, and its injection into early *Xenopus* embryos caused craniofacial abnormalities such as small and deformed heads, malformed auditory organs, and abnormal or misplaced eyes. Ductin-independent reagents that affect membrane voltage produced similar results, so the abnormalities were not caused directly by the mutant protein. The voltage-reporting dyes CC2-DMPE and DiBAC₄(3) confirmed that the treatments had altered the bioelectrical patterns of the embryos, and in situ hybridization showed that the altered bioelectrical patterns corresponded to disrupted expression patterns of several genes involved in craniofacial development [277].

Vaibhav Pai, Michael Levin and their colleagues also manipulated ion channels in *Xenopus* embryos to induce changes in transmembrane voltage potential that disrupted eye patterning. Using CC2-DMPE, they first showed that eyes normally develop from two small areas in the anterior neural field that are hyperpolarized (internally more negative, by about 10mV) relative to neighboring tissues. To determine whether hyperpolarization is required for normal eye development, the researchers injected the dorsal blastomeres of four-cell-stage *Xenopus* embryos with mRNAs for two ion channels that are not normally present: EXP1, which conducts positively charged ions into the cell [278], and GlyR, which can be chemically activated to enable the outflow of Cl⁻ ions [279,280]. Since hyperpolarization requires the exclusion of positively charged ions and the retention of negatively charged ions, these two channels strongly disrupted the hyperpolarization patterns in injected embryos. The resulting tadpoles had a high incidence of incomplete, small, or deformed eyes—or they were missing eyes entirely. Control embryos that were uninjected, injected with a non-channel protein, or injected with GlyR alone or its chemical activator alone, developed well-formed eyes. Various other ion channels were also used to perturb the normal pattern of transmembrane voltage potential. The results included not only the eye defects listed above, but also ectopic eyes well outside the anterior neural field—including eyes on the tadpole's side or tail [281].

EEFs as spatial coordinate systems

In the 1980s, Jaffe and Nuccitelli both proposed that ionic currents and/or voltage gradients could provide ontogenetic information for embryonic pattern formation, though they acknowledged that many experiments remained to be done [248,282]. Experiments such as those described above clearly demonstrate that the disruption of bioelectric patterns has significant local effects on ontogeny, but EEFs might play a larger role as well.

Harold Burr and Filmer Northrop suggested in the 1930s that endogenous “electro-dynamic fields” determine (or at least strongly influence) the global anatomy of an organism [220]. In 1995, Shi and Borgens reviewed the experimental evidence that embryonic cells are responsive to applied voltages in the physiological range and that disruption of EEFs produces developmental defects consistent with the disruption. They concluded that EEFs “may provide a three dimensional coordinate system” that helps to specify form in embryos [283].

The term “morphogenetic field” has had many meanings since the idea of embryonic fields was first proposed by Alexander Gurwitsch in the early twentieth century [284–286]. As used by Michael Levin, the field concept denotes both informational and regional relationships, and its essence is “non-locality”—the idea that “many diverse examples of pattern formation are best understood not as cell-level behaviors around any one locale but rather at higher levels of organization.” Although gradients of morphogenetic molecules may specify some patterns, Levin (following Burr and others) argues that EEFs may also function as “templates of shape.” Thus “it is likely that a full understanding of the morphogenetic field and its informational content will need to involve cracking the bioelectric code (the mapping between spatiotemporal ionic profile patterns and tissue morphology outcomes)” [287,288].

MEMBRANE PATTERNS AND DNA

So membrane patterns—the three-dimensional arrangements of membrane-associated proteins, lipids and carbohydrates, as they change over time—carry essential ontogenetic information. Yet (as I demonstrate below) the information carried by membrane patterns cannot be reduced to sequence information in DNA, for at least two reasons. First, the vast majority of proteins in eukaryotes are not completely specified by DNA sequences. Second, even if DNA sequences completely specified all proteins, DNA would not specify their spatiotemporal arrangements in membranes.

Most proteins are not completely specified by DNA sequences

The central dogma (which here includes Crick's sequence hypothesis) claims that (1) DNA specifies RNA and (2) RNA specifies protein. Yet this claim fails at both steps, because most RNAs are not uniquely specified by DNA sequences, and many proteins are not uniquely specified by RNAs—either in their amino acid sequences or in their final folded forms.

After transcription, RNAs from many eukaryotic genes undergo alternative splicing. Recent studies estimate that transcripts from approximately 95% of multi-exon human genes are spliced in more than one way [289–291]. By intervening between transcription and translation, alternative splicing generates RNAs with sequences that differ from DNA sequences [292]. The differences are functionally significant. In humans and other animals, alternatively spliced transcripts are expressed in tissue- and developmental stage-specific ways [293,294]. Among other things, they regulate physiological

changes [295], neuronal development [296], and stem cell pluripotency [297]. Alternative splicing is essential for the transition between epithelial cells and mesenchymal cells [298,299], and it produces enormous diversity in membrane proteins, including ion channels [300,301]. With alternative RNA splicing, cells make thousands more RNAs and proteins than are encoded in DNA sequences. One immunoglobulin gene in *Drosophila melanogaster* generates more than 18,000 protein isoforms through alternative splicing [302].

In addition to alternative splicing, many metazoan transcripts undergo RNA editing, which can (a) modify cytidine to uridine; (b) modify adenosine to inosine; or (c) insert additional nucleotides. Several recent analyses have demonstrated extensive RNA editing in the human transcriptome [303–305]. The editing of an mRNA often alters the amino acid sequence of the encoded protein so that it differs from the sequence predicted by the DNA [306,307]. RNA editing can also regulate the amount of functional RNA in the cell and expand the functionality of a limited set of transcripts [308]. One example of functional editing is found in two species of octopus, one living in the Antarctic and the other in the Caribbean. Both possess very similar genes for a particular potassium channel, but in the Antarctic species RNA editing recodes an isoleucine to a valine in the channel's pore, greatly accelerating its gating kinetics so that it can function properly in the extreme cold [309]. Other examples of functional RNA editing involve adenosine deaminases that act on RNAs (ADARs), which selectively convert adenosine (A) to inosine (I) [310]. ADARs are widespread in metazoan nervous systems, where they edit pre-mRNAs of proteins involved in electrical and chemical neurotransmission [311–314]. In mice, ADARs are necessary for the survival of the stem cells that generate mature blood cells [315], and they play an important role in the gastrointestinal tract [316].

Not only are the nucleotide sequences of many RNAs not fully specified by DNA sequences, but also the amino acid sequences and final folded forms of many proteins—on which their functions depend—are not fully specified by RNA sequences. Some proteins undergo splicing, a phenomenon first reported in yeast in 1990 [317,318]. In protein splicing, an “intein” (analogous to an intron in RNA splicing) is removed from the protein and the flanking pieces (“exteins”) are spliced together without the need for cofactors or enzymes [319–323]. Although protein splicing has been found only in unicellular organisms, a somewhat analogous process occurs in the biogenesis of various metazoan proteins. The process in metazoans is called “autoproteolysis,” in which a protein cleaves itself into two pieces. Autoproteolysis is widespread in the hedgehog family of proteins (needed for signal transduction in animal development) and in nuclear pore complex proteins [324–328].

Even in many proteins that do not undergo splicing, the amino acid sequence does not completely specify the final folded form. Protein folding can be affected by the intracellular milieu, including factors such as pH, the variety of molecular chaperones, and the degree of macromolecular crowding

[329–334]. Some proteins adopt similar forms and/or functions despite having very dissimilar amino acid sequences [335–339]. Others assume different three-dimensional forms despite having the same or very similar amino acid sequences. Some well-known examples of the latter are prions—misfolded proteins that are pathogenic and heritable [340–343]. But there are also non-pathogenic proteins in which the same amino acid sequence can fold into more than one shape; these are known as “metamorphic” proteins [344,345]. One example of a metamorphic protein is Mad2 (for “mitotic arrest deficient”), a protein that monitors the attachment of microtubules to kinetochores [346,347]. Another example is lymphotactin, a small secreted signaling molecule [348,349]. A third example is the CLIC family of proteins, which function in chloride channels [350,351]. Still another example is IscU, which is involved in iron-sulfur cluster assembly in bacteria and eukaryotic mitochondria [352,353].

Some proteins are “intrinsically disordered.” Approximately 20–30% of proteins in mammalian cells are inherently devoid of any ordered three-dimensional structure and adopt folded conformations only after interacting with other molecules [354]. More than 50% of eukaryotic proteins have at least one domain longer than 30 amino acids that is intrinsically disordered [355–358]. Intrinsically disordered proteins play crucial roles in regulating ion channels and serving as molecular hubs in intracellular signaling networks [359–361].

Finally, most eukaryotic proteins are post-translationally modified by glycosylation [362]. First a glycan is attached to the protein, usually through an asparagine, serine or threonine residue [363–366]. In some cases the residue is part of a consensus sequence, but in other cases it is not [367,368]. For branched glycoproteins, enzymatic networks then add or remove carbohydrate residues in tissue- and developmental stage-specific patterns [369,370]. For unbranched proteoglycans, other enzymatic networks vary the order and sulfation patterns of the constituent disaccharides to yield different spatial and temporal distributions [371,372]. These enzymatic networks are quite complex, involving dozens of interacting glycotransferases, and they can vary glycosylation patterns epigenetically depending on the needs of the cell in a given environment or at a specific developmental stage [373–375]. The final glycosylated form of a protein is thus very far removed from direct specification by a DNA sequence. As various biochemists have put it, the modification of proteins through enzymatic glycosylation imparts “an additional level of ‘information content’ to underlying polypeptide structures” [376]; it “is an event that reaches beyond the genome” [377]; and if DNA-centered biology is compared to Newtonian physics, protein glycosylation may pave the way for a “quantum mechanics of biology” and a “scientific revolution analogous to the one which transformed the field of physics in the early 20th century” [378].

Membrane patterns are not specified by DNA

On the second point, even if DNA sequences uniquely specified the molecular structures of proteins, DNA would

not specify the spatial distribution of proteins in the plasma membrane. Some membrane patterns are templated by the membranes from which they are derived, with proteins from the cell interior being incorporated during membrane growth only if they match the existing matrix [379]. Such templating has been well studied in protozoa. In 1965, Janine Beisson and Tracy Sonneborn induced one member of a conjugating pair of *Paramecium aurelia* to transfer to its partner a section of cortex that had been surgically inverted 180° relative to the surrounding cortex. Although the DNA was unchanged, the altered pattern of cilia in the cortex was heritable. Indeed, ciliates with artificially inverted rows have been stably maintained for thousands of generations [380–382]. In 1977, Stephen Ng and Joseph Frankel observed the same phenomenon in *Tetrahymena pyriformis* and concluded, “The cell as an architect thus not only makes use of the genomic information to produce the appropriate building blocks, but, in addition, also arranges the building blocks according to the blueprint as defined in the preexisting architecture” [383]. Frankel called this extra-genic blueprint the “corticotype” [384]. Similar results were reported in *Tetrahymena pyriformis* by David Nanney and in *Stylonychia mytilus* by Gary Grimes [385,386].

Cortical templating is not limited to protozoa. In 1930, Sonneborn reported it in a metazoan, the flatworm *Stenostomum incaudatum* [387,388]. In 1990, Michael Locke observed paired patterns in caterpillar epidermis cells that “imply that a part of the epigenetic sequence leading to the formation of the pattern has replicated [and been] inherited by daughter cells. It is not just genetic material that is inherited but part of a cell in a particular state.” According to Locke, such inheritance “requires more than number and kind of molecule. The duplication of pattern involves relative position and orientation,” factors that “cannot be specified only by a base sequence” [389].

In 1983, Robert Poyton proposed that biological membranes carry “spatial memory,” the units of which are spatially localized hetero-oligomeric proteins. According to Poyton, when phospholipids are incorporated into a growing membrane in preparation for replication, the hetero-oligomers dissociate into their subunits. Newly synthesized subunits in the cytoplasm then associate with the corresponding older subunits to form hybrid hetero-oligomers that are chemically identical to the originals. Thus each unit of spatial memory would replicate semi-conservatively, like DNA (Fig. 8) [390]. Consistent with Poyton’s hypothesis is evidence that local lipid bilayer properties influence the stability of transmembrane oligomers [391–393]; that protein dimers in membranes dissociate when diluted with phospholipids [394,395]; that membrane proteins selectively recruit other proteins in ESCRT complexes and Ras nanoclusters [396–401]; and that intracellular membranes grow by incorporating new proteins into pre-existing templates [402]. Poyton concluded, “Realizing that genetic memory is one-dimensional, along a DNA molecule, whereas spatial memory is likely to be two-dimensional, along membrane surfaces, and three-dimensional within the cellular interior, it is probable that spatial memory is more complicated and diverse than genetic memory” [390].




Figure 8. Membrane replication. In Robert Poyton’s theory of membrane replication, the unit of membrane memory is a hetero-oligomeric protein, represented here in simplified form by the two rectangles (A). As the membrane grows from the addition of new phospholipids, the parts of the oligomer separate (B). Newly synthesized proteins from the cytoplasm associate with the complementary subunits from the original oligomer (C). The result is two new oligomers that are chemically identical to the originals (D). doi:10.5048/BIO-C.2014.2.f8

A eukaryotic cell contains many different types of membranes, which Thomas Cavalier-Smith has collectively designated the cell’s “membranome” [403]. Each membrane is characterized not only by its chemistry but also by its topology. Some recent writers have used “membranome” in a reductionist sense to mean the list of individual proteins found in membranes, though this is clearly at odds with Cavalier-Smith’s original topological formulation [404–406].

Cavalier-Smith distinguished between “genetic membranes” and “derived membranes.” Genetic membranes, which include the endoplasmic reticulum and plasma membranes of many cells, “always arise by growth and division of the same type.” Derived membranes, which include lysosomal membranes, “form instead by differentiation from dissimilar membranes.” Some genetic membranes are “primary,” that is, they grow by direct insertion of individual protein and lipid molecules. Other genetic membranes are “secondary,” in the sense that they grow (in eukaryotes) by incorporation of vesicles from a primary genetic membrane [407]. If any type of genetic membrane were lost, it could probably not be regenerated from its constituent molecules—even if all the genes encoding its proteins and lipid-synthesizing machinery remained—because the requisite spatial pattern would be gone.

The closest thing to an exception to this last statement is found in *Bryopsis*, a genus of green algae consisting of large multinucleate cells. When one of these cells is damaged or fragmented it forms protoplasts in seawater that are not enclosed in plasma membranes, but in gelatinous envelopes made of polysaccharides. After about 12 hours a lipid bilayer might form underneath the polysaccharide coat—but only if a protoplast retains at least 15% of the original cell membrane. If such a protoplast also contains a nucleus, it might regenerate into a complete plant [408–413]. So even in the case of *Bryopsis* protoplasts, new plasma membrane is regenerated “by growth and division of the same type” of membrane, as Cavalier-Smith pointed out. It does not form de novo. Furthermore, the

membrane regeneration seen in Bryopsis appears to be unique; it is not found in other plants, in bacteria, in protozoa, or in metazoa—in which membrane patterns carry more information than can be preserved in a small fragment.

The importance of global spatial information can be illustrated by the first cell division of a *Xenopus* zygote. The animal-vegetal axis and cortical rotation divide the cortex into four broad zones. In Figure 9, the intersection of planes A and B corresponds to the animal-vegetal axis, while the intersection of planes A and C corresponds to the direction of cortical rotation. If the zygote divides along plane A, each of the two daughter cells will inherit not only a nucleus but also portions of all four zones of cortical information, so if the daughter cells are separated they can both develop into complete tadpoles. If the zygote divides along plane B or C, however, each daughter cell will inherit a nucleus, but not a full set of cortical information, so they cannot develop into two tadpoles.

Thus, according to Cavalier-Smith, the idea that the genome contains all the information needed to make an organism “is simply false. Membrane heredity, by providing chemically specific two-dimensional surfaces with mutually conserved topological relationships in the three spatial dimensions, plays a key role in the mechanisms that convert the linear information of DNA into the three-dimensional shapes of single cells and multicellular organisms. Animal development creates a complex three-dimensional multicellular organism not by starting from the linear information in DNA... but always starting from an already highly complex three-dimensional unicellular organism, the fertilized egg, which membrane and DNA heredity together have perpetuated” [403].




Figure 9: Zones of cortical information in a *Xenopus* zygote. The animal-vegetal axis and cortical rotation divide the cortex of a *Xenopus* zygote into four broad zones. Here the intersection of planes A and B corresponds to the animal-vegetal axis, while the intersection of planes A and C corresponds to the direction of cortical rotation. If the zygote (after duplication of the nucleus) divides along plane A, each of the two daughter cells will inherit not only a nucleus but also portions of all four zones of cortical information; if these daughter cells are separated, they can both develop into complete frogs. If the zygote divides along plane B or C, however, each daughter cell will inherit a nucleus, but not a full set of cortical information; their development will be blocked.

doi:10.5048/BIO-C.2014.2.f9

Membrane heredity, however, cannot explain how membranes change in specified ways during ontogeny. In most cases, embryonic cells do not simply transmit duplicate membrane patterns to their daughter cells when they divide. The membrane patterns of nerve cells are very different from those of muscle cells, blood cells, bone cells, and liver cells—to name only a few—so membrane patterns have to change in the course of differentiation and morphogenesis. The changes are not haphazard, however; they are highly regulated to generate the integrated organ systems of an adult of the proper species. If membrane patterns are not specified by DNA sequences or by patterns in the cells from which they are derived, how are they specified? I will return to this question below.

SUMMARY AND IMPLICATIONS

In 1958, Francis Crick proposed that the specificity of a DNA segment lies solely in its nucleotide sequence, which encodes the nucleotide sequence of a messenger RNA (mRNA) and thereby the amino acid sequence of a protein. Crick also proposed that sequence information can be transferred from DNA to RNA to protein, but not in the other direction [1]. Crick called these the “sequence hypothesis” and the “central dogma,” respectively. For brevity (and in accordance with common usage), I subsume both under the title “central dogma,” which has been popularly stated as “DNA makes RNA makes protein makes us.”

The central dogma logically implies that embryo development (ontogeny) is directed by a genetic program encoded in DNA sequences. Elaborating on this, Eric Davidson maintains that developmental gene regulatory networks (dGRNs) “control ontogeny of the body plan” [15]. But dGRNs must be distributed in embryonic space, and (as Davidson acknowledges) their distribution depends on spatial anisotropies in the egg that are defined in relation to the embryonic axes that establish the basic body plan.

Yet the evidence from oogenesis in *Drosophila*, *Xenopus*, and echinoderms shows that the major embryonic axes are established long before the embryo’s dGRNs are put in place—much less activated. The anterior-posterior axis in *Drosophila*, the animal-vegetal axis in *Xenopus*, and the animal-vegetal axis in echinoderms are initially derived from the architecture of the ovary, through processes mediated by cytoskeletal and membrane patterns rather than dGRNs. Some might argue that the architecture of the ovary was set up by GRNs in the mother, but in the absence of sufficient evidence such an argument is simply a restatement of the central dogma. In embryo development, there is no empirical justification for giving DNA sequences ontogenetic priority over architectural constraints and cytoskeletal and membrane patterns.

Although cytoskeletal and membrane patterns interact in complex ways and cannot be completely understood apart from each other, this review has focused primarily on the latter. Biological membranes were once thought to be like two-dimensional seas in which proteins float and diffuse freely, unconstrained except for local interactions. It is now clear, however, that many membrane proteins and lipids are compartmentalized into spatially restricted domains and

arranged in global patterns that affect ontogeny.

Some plasma membrane patterns are formed in response to extracellular signals, as is the case with mesenchymal and epithelial cells; others are less dependent on the extracellular environment and more intrinsic to the cell, as is the case with spatially localized glycans and ion channels. Plasma membrane patterns serve essential ontogenetic functions by providing targets and sources for intracellular signaling and transport, by regulating cell-cell interactions, and by generating endogenous electric fields that provide three-dimensional coordinate systems for embryo development.

Membrane patterns are not specified by DNA sequences. First, DNA sequences only partially specify RNAs and proteins. After transcription, many RNAs undergo alternative splicing and/or editing, so thousands of different mRNAs can be generated from a single DNA sequence. After translation, some proteins are edited to produce different amino acid sequences, and many proteins with similar amino acid sequences can adopt more than one folded structure—or they are intrinsically disordered. Furthermore, most eukaryotic proteins are post-translationally modified by glycosylation. Given the enormous number of possible glycan structures, a protein can be modified in trillions of possible ways. If “makes” is taken to mean “specifies,” then “DNA makes RNA makes protein” fails at each step.

Second, even if DNA completely specified proteins, it would not follow that one-dimensional DNA sequences specify the two- and three-dimensional arrangements of proteins in membranes. Many membrane patterns are templated by the membranes from which they are derived, and proteins from the cell interior are incorporated during membrane growth only if they match the existing matrix. Such templating has been especially well studied in protozoa, but it has also been observed in planaria and insect epidermis. According to evolutionary biologist Thomas Cavalier-Smith, the idea that the genome contains all the information needed to make an organism “is simply false. Membrane heredity, by providing chemically specific two-dimensional surfaces with mutually conserved topological relationships in the three spatial dimensions, plays a key role in the mechanisms that convert the linear information of DNA into the three-dimensional shapes of single cells and multicellular organisms” [403].

In embryo development, however, membrane heredity cannot be the whole story. During ontogeny many new membrane patterns arise that cannot be traced back to patterns in pre-existing membranes. The new patterns do not arise haphazardly; they are highly specified. Yet there is no evidence that they—any more than the patterns that precede them—are determined by a program in the organism’s DNA. Whether membrane patterns are templated or form *de novo*, they carry ontogenetic information that is specified independently of DNA sequences [414]. This fact has serious implications both for evolutionary theory and for our understanding of ontogeny.

Implications for evolutionary theory

Of course, there is no single theory of evolution. First, the word “evolution” has many meanings, including simple

change over time, the history of the cosmos, or (in biology) the transformation of one species into another. Second, even in biology there are several theories of evolution. I will focus on one of these, which I will call Neo-Darwinism.

Like evolution itself, Neo-Darwinism has had many meanings. The term was first used by Samuel Butler in 1880 to distinguish Charles Darwin’s theory from that of his grandfather, Erasmus Darwin. Butler used “Darwinism” to refer approvingly to Erasmus’s theory that new variations arise “due to the wants and endeavours of the living forms in which they appear,” while Butler used “Neo-Darwinism” to refer disparagingly to Charles’s ascription of new variations “to chance, or, in other words, to unknown causes” [415,416].

In 1895, Georges Romanes used “Neo-Darwinian” to describe the view (which he attributed to August Weismann and Alfred Russel Wallace) that “natural selection is the only possible cause of adaptive modification;” Romanes used “Neo-Lamarckian” to describe the view (which he attributed mainly to Americans) that “much greater importance ought to be assigned to the inherited effects of use and disuse than was assigned to these agencies by Darwin.” According to Romanes, Charles Darwin’s view (which he called “Darwinism”) stood “between these two extremes” [417].

When Charles Darwin’s theory of evolution was combined in the 1930s with Gregor Mendel’s theory of genetics, the resulting “modern synthesis” attributed new variations to genetic mutations and excluded the inherited effects of use and disuse [418,419]. Since then, many people have used the term Neo-Darwinism to refer to the modern synthesis, as supplemented by molecular genetics.

I use Neo-Darwinism here to mean the view that all living things are descendants of one or a few ancestral organisms that have been modified principally (though not exclusively) by the natural selection of advantageous variations, and that new variations originate from changes in DNA sequences. This was Jacques Monod’s view, though he called it Darwinism [11]. Eric Davidson distances himself from what he calls the “neo-Darwinian concept” that evolution occurs by “small continuous changes in primary protein sequence,” and he attributes evolution instead to system level changes in developmental gene regulatory networks (dGRNs) [15]. But I would argue that in so far as he attributes control of ontogeny to a program in DNA, Davidson’s view is still a variant of Neo-Darwinism.

As we have seen, however, the idea that embryo development is controlled by a genetic program is inconsistent with the biological evidence. Embryo development requires far more ontogenetic information than is carried by DNA sequences. Thus Neo-Darwinism is false.

Yet population genetics, which assumes that evolution is due to changes in gene frequencies, is based on the assumption that Neo-Darwinism is true. As originally formulated in the 1920s and 1930s by J. B. S. Haldane, Ronald Fisher, and Sewall Wright, population genetics relied on Mendelian genes [421–423], while modern population genetics assumes that those genes are equivalent to DNA sequences. A recent advocate of the centrality of population genetics is Michael Lynch, who

wrote in 2006 that “evolution is a population genetic process governed by four fundamental forces.” One force is selection, “whose central role in the evolution of complex phenotypic traits is universally accepted, and for which an elaborate formal theory in terms of genotypic frequencies now exists.” The other three forces are mutation (including DNA insertions, deletions and duplications), recombination, and random genetic drift. “Given the century of empirical and theoretical work devoted to the study of evolution,” Lynch wrote, “the only logical conclusion is that these four broad classes of mechanisms are, in fact, the only fundamental forces of evolution.” Thus “nothing in evolution makes sense except in the light of population genetics” [424].

Of course, no one denies that DNA is essential for ontogeny, and that DNA mutations can affect phenotypic traits. Furthermore, population genetics may be useful for tracing lineages within existing species. But since the ontogenetic information in an embryo far exceeds that in its DNA, evolution must necessarily involve far more than changes in gene frequencies.

In 1992, Brian Hall noted that population genetics is “neither sufficient nor inclusive” as an explanation for evolution. He proposed the term “evolutionary developmental biology” (which others had used before him) to describe the study of “how developmental processes effect evolutionary change and how development itself has evolved.” Hall concluded: “Evolution acts at the three levels of changes in gene frequencies, the appearance of new characters, and the adaptation and radiation of new species. The common denominator of all three is genetic change through time, the common agent of that change is alteration in ontogeny, the common integrator of the three is epigenetic organization. The science of the study of these interactions is evolutionary developmental biology” [425].

Despite Hall’s inclusion of ontogeny and epigenetic organization, many advocates of “evo–devo” (the nickname for evolutionary developmental biology) have perpetuated the Neo-Darwinian emphasis on genetic programs. For example, Sean B. Carroll wrote in 2005 that evo–devo focuses on “tool-kit genes”—developmental genes that are common to a wide variety of multicellular animals but controlled by different genetic switches “that integrate inputs from other tool-kit proteins acting a bit earlier in the embryo.” The embryo has a coordinate system that “imposes some spatial order on how the program of tool-kit genes unfolds,” but Carroll emphasized the primacy of interacting genes and gene products: “The important point to know is that the throwing of every switch is set up by preceding events, and that a switch, by turning on its gene in a new pattern, in turn sets up the next set of patterns and events in development.” Thus the rules for generating animal form are found in DNA: “In the entire complement of DNA of a species (the genome), there exists the information for building that animal.” The “evolution of form is ultimately then a question of genetics” [426]. So I would argue that Carroll’s view (like Davidson’s) is a variant of Neo-Darwinism, and thus false.

An adequate theory of evolution would not try to force organisms into the Procrustean bed of the central

dogma—though basing a theory of evolution solely on changes in membrane patterns would be equally mistaken. The latter carry ontogenetic information that is specified independently of DNA sequences, but a case could be made (though I have not made it here) that the same is true of cytoskeletal patterns. So an adequate theory of evolution would have to explain how various information sources in the organism (including its DNA, membrane patterns, and cytoskeletal patterns) change in a coordinated fashion to produce new species, organs, and body plans. Before attempting to explain how organisms change in the course of phylogeny, however, we need to address the question of how they change in the course of ontogeny.

Implications for our understanding of ontogeny

A developing animal embryo undergoes thousands of changes in its molecules, cells, organs, and overall anatomy. Such changes are integrated to produce, with great fidelity, a specific outcome. Although disrupting development can produce death or deformity, embryos have a remarkable ability to recover from disruptions and produce normal adults of their species. Ontogeny is thus orchestrated, reliable, robust, and stereotypical. What is the best way to understand it?

One way is molecular biology, which takes an essentially Newtonian approach. Living systems are analyzed in terms of molecules (analogous to atoms in Newtonian physics), their locations and movements in the cell (position and velocity in Newtonian mechanics), and their chemical and physical interactions (the forces among them). Given an embryo in its initial state, all subsequent states (including new membrane patterns) would follow from the ordinary laws of physics acting on the constituent molecules. This does not preclude the emergence of unexpectedly complex and dynamic patterns, analogous to those we see in the Belousov-Zhabotinsky reaction [427,428], but such patterns would be only “weakly” emergent: Though unexpected, they arise from the lower-level constituents of a system by the ordinary operation of the laws of classical (and perhaps quantum) mechanics [429,430].

But what if embryo development involves “strongly” emergent properties? What if ontogeny requires higher-level information that cannot be reduced to the characteristics and interactions of lower-level constituents [431]? In a 2012 discussion of the bioelectric code, Michael Levin proposed that “biological structures encode maps for their shape”—what he called “target morphologies.” According to Levin,

What makes target morphology models distinct from [weakly] emergent models is the hypothesis that there are some measurable quantities contained in the living system that are directly isomorphic to the anatomy that is being constructed or maintained. In [weakly] emergent models, there is no such process, the shape being assembled as the result of low-level rules and not by comparison to (or directives from) any informational structure that encodes a final shape” [287].

A Newtonian approach permits weakly emergent properties

but excludes strongly emergent ones; yet it is not the only way to approach living systems scientifically. In the 1950s, mathematical biologists Nicholas Rashevsky and Robert Rosen introduced a new approach they called “relational biology” [432–434]. Unlike Newtonian biology, which gives ontological priority to matter (i.e., molecules), relational biology (as the name implies) gives ontological priority to the relations that constitute an organized system. Although molecular biology has been successful at a certain level, its methods involve discarding the organization of a cell while keeping the matter. Yet the former, once discarded, cannot be recovered from the latter—and living things are fundamentally characterized by their organization.

Relational biology has radical implications not only for biology, but also for physics. A Newtonian approach assumes that living things embody a subset of the natural laws governing atoms and the universe; thus the former are in principle reducible to the latter. But if living things embody natural laws in addition to those that govern non-living matter, the Newtonian approach is mistaken. Relational biology begins with the opposite assumption—namely, that atoms and the universe embody a subset of the natural laws governing living things. The latter can then be studied in their own right, without trying to reduce them to the former [435]. Of course, to be scientifically fruitful relational biology must move beyond mere assumption.

In the 1940s, seventy years after the formulation of classical set theory, Samuel Eilenberg and Saunders Mac Lane proposed a more general theory of structures and systems of structures called “category theory,” which was further developed in the 1950s and 1960s by Alexander Grothendieck, Peter Freyd and William Lawvere [436–440]. A category consists of “objects”

and “mappings.” A category-theoretic mapping (also called a transformation or morphism) is a process for getting from one object to another object; it is customarily represented by an arrow. A category-theoretic object is a vertex from which one or more arrows originate or at which one or more arrows terminate. It is a logico-mathematical entity (such as a set), not necessarily a physical thing (though a set can consist of one or more physical things). Category theory is primarily concerned with the analysis of mappings and their interrelationships. In set theory, a “function” is a mapping from an element of one set to an element of another set; in category theory, a “functor” is a mapping from one category to another that preserves the relational structures of the mappings and objects. Since 1945, category theory has been logically and mathematically worked out in great detail [441–443].

Category theory provides a rigorous conceptual foundation for relational biology. Starting with Rosen in 1958, biologists have now applied it to various aspects of living systems [444–448]. Much more theoretical and experimental work remains to be done, of course. One promising research program will involve applying relational biology and category theory to the DNA-independent specification of ontogenetic information in membrane patterns.

Acknowledgements

The author gratefully acknowledges valuable discussions with Richard von Sternberg, and he thanks Russell W. Carlson, Paul A. Nelson, Richard Nuccitelli, and Jason Tresser for critically reading sections of the manuscript. The author also gratefully acknowledges helpful comments from several anonymous reviewers. Marco Doelling produced the illustrations. Funding was provided by the Discovery Institute, Seattle, Washington.

1. Crick FHC (1958) On protein synthesis. *Symp Soc Exp Biol* 12:138-163.
2. Jacob F, Monod J (1961) Genetic regulatory mechanisms in the synthesis of proteins. *J Mol Biol* 3:318-356. doi:10.1016/S0022-2836(61)80072-7
3. Jacob F (2011) The birth of the operon. *Science* 332:767. doi:10.1126/science.1207943
4. Davidson EH (1968) *Gene Activity in Early Development*. Academic Press (New York) pp 3,8,12,154.
5. Sonneborn TM (1970) Gene action in development. *Proc R Soc Lond B* 176:347-366. doi:10.1098/rspb.1970.0054
6. Sapp J (1987) *Beyond the Gene: Cytoplasmic Inheritance and the Struggle for Authority in Genetics*. Oxford University Press (New York).
7. Jacob F (1970) *La Logique du Vivant*. Editions Gallimard (Paris). Translated by Spillmann BE (1973) as *The Logic of Life*. Princeton University Press (Princeton) p 3.
8. Monod J (1970) *Le Hasard et la Necessité*. Editions du Seuil (Paris). Translated by Autry Wainhouse (1971) as *Chance and Necessity*. Alfred A. Knopf (New York) p 109.
9. Keller EF (2000) *The Century of the Gene*. Harvard University Press (Cambridge, MA).
10. Sapp J (2003) *Genesis: The Evolution of Biology*. Oxford University Press (New York).
11. Judson HF (1979) *The Eighth Day of Creation: The Makers of the Revolution in Biology*. Simon and Schuster (New York) p 217.
12. Kauffman SA (1969) Metabolic stability and epigenesis in randomly constructed genetic nets. *J Theor Biol* 22:437-467. doi:10.1016/0022-5193(69)90015-0
13. Davidson EH (1990) How embryos work: A comparative view of diverse modes of cell fate specification. *Development* 108:365-389.
14. Davidson EH (2006) *The Regulatory Genome: Gene Regulatory Networks in Development and Evolution*. Academic Press (Amsterdam) p 94.
15. Davidson EH (2011) Evolutionary bioscience as regulatory systems biology. *Dev Biol* 357:35-40. doi:10.1016/j.ydbio.2011.02.004
16. Nüsslein-Volhard C, Frohnhöfer HG, Lehmann R (1987) Determination of anteroposterior polarity in *Drosophila*. *Science* 238:1675-1681. doi:10.1126/science.3686007
17. Anderson KV, Nüsslein-Volhard C (1984) Information for the dorsal-ventral pattern of the *Drosophila* embryo is stored in maternal RNA. *Nature* 311:223-227. doi:10.1038/311223a0
18. Schüpbach T, Wieschaus E (1986) Germline autonomy of maternal-effect mutations altering the embryonic body pattern of *Drosophila*. *Dev Biol* 113:443-448. doi:10.1016/0012-1606(86)90179-X

19. Driever W, Nüsslein-Volhard C (1988) The *bicoid* protein determines position in the *Drosophila* embryo in a concentration-dependent manner. *Cell* 54:95-104. doi:10.1016/0092-8674(88)90183-3
20. Lehmann R, Nüsslein-Volhard C (1991) The maternal gene *nanos* has a central role in posterior pattern formation of the *Drosophila* embryo. *Development* 112:679-691.
21. Wharton RP, Struhl G (1991) RNA regulatory elements mediate control of *Drosophila* body pattern by the posterior morphogen *nanos*. *Cell* 67:955-967. doi:10.1016/0092-8674(91)90368-9
22. Lewis EB (1978) A gene complex controlling segmentation in *Drosophila*. *Nature* 276:565-570. doi:10.1038/276565a0
23. Nüsslein-Volhard C, Wieschaus E (1980) Mutations affecting segment number and polarity in *Drosophila*. *Nature* 287:795-801. doi:10.1038/287795a0
24. Lawrence PA, Morata G (1994) Homeobox genes: Their function in *Drosophila* segmentation and pattern formation. *Cell* 78:181-189. doi:10.1016/0092-8674(94)90289-5
25. Carroll SB (2008) Evo-devo and an expanding evolutionary synthesis: A genetic theory of morphological evolution. *Cell* 134:25-36. doi:10.1016/j.cell.2008.06.030
26. Spirov A, Fahmy K, Schneider M, Frei E, Noll M, et al. (2009) Formation of the *bicoid* morphogen gradient: An mRNA gradient dictates the protein gradient. *Development* 136:605-614. doi:10.1242/dev.031195
27. Lipshitz HD (2009) Follow the mRNA: A new model for *Bicoid* gradient formation. *Nat Rev Mol Cell Biol* 10:509-512. doi:10.1038/nrm2730
28. Theurkauf WE, Hazelrigg TI (1998) In vivo analyses of cytoplasmic transport and cytoskeletal organization during *Drosophila* oogenesis: Characterization of a multi-step anterior localization pathway. *Development* 125:3655-3666.
29. St Johnston D (2005) Moving messages: The intracellular localization of mRNAs. *Nat Rev Mol Cell Biol* 6:363-375. doi:10.1038/nrm1643
30. Weil TT, Xanthakis D, Parton R, Dobbie I, Rabouille C, et al. (2010) Distinguishing direct from indirect roles for *bicoid* mRNA localization factors. *Development* 137:169-76. doi:10.1242/dev.044867
31. Sardet C, Prodon F, Dumollard R, Chang P, Chênevert J (2002) Structure and function of the egg cortex from oogenesis through fertilization. *Dev Biol* 241:1-23. doi:10.1006/dbio.2001.0474
32. Irion U, St Johnston D (2007) *bicoid* RNA localization requires specific binding of an endosomal sorting complex. *Nature* 445:554-558. doi:10.1038/nature05503
33. Lasko P (2012) mRNA localization and translational control in *Drosophila* oogenesis. *Cold Spring Harb Perspect Biol* 4:a012294. doi:10.1101/cshperspect.a012294
34. González-Reyes A, St Johnston D (1994) Role of oocyte position in establishment of anterior-posterior polarity in *Drosophila*. *Science* 266:639-642. doi:10.1126/science.7939717
35. Lin H, Yue L, Spradling AC (1994) The *Drosophila* fusome, a germline-specific organelle, contains membrane skeletal proteins and functions in cyst formation. *Development* 120:947-956.
36. Huynh J-R, St Johnston D (2004) The origin of asymmetry: Early polarisation of the *Drosophila* germline cyst and oocyte. *Curr Biol* 14:R438-R449. doi:10.1016/j.cub.2004.05.040
37. Grieder NC, de Cuevas M, Spradling AC (2000) The fusome organizes the microtubule network during oocyte differentiation in *Drosophila*. *Development* 127:4253-4264.
38. Bolívar J, Huynh JR, López-Schier H, González C, St Johnston D, et al. (2001) Centrosome migration into the *Drosophila* oocyte is independent of BicD and egl, and of the organisation of the microtubule cytoskeleton. *Development* 128:1889-1897.
39. Ray RP, Schüpbach T (1996) Intercellular signaling and the polarization of body axes during *Drosophila* oogenesis. *Genes Dev* 10:1711-1723. doi:10.1101/gad.10.14.1711
40. Bilder D, Haigo SL (2012) Expanding the morphogenetic repertoire: Perspectives from the *Drosophila* egg. *Dev Cell* 22:12-23. doi:10.1016/j.devcel.2011.12.003
41. Roth S, Lynch JA (2009) Symmetry breaking during *Drosophila* oogenesis. *Cold Spring Harb Perspect Biol* 1:a001891. doi:10.1101/cshperspect.a001891
42. Vincent J-P, Oster GE, Gerhart J (1986) Kinematics of grey crescent formation in *Xenopus* eggs: The displacement of subcortical cytoplasm relative to the egg surface. *Dev Biol* 113:484-500. doi:10.1016/0012-1606(86)90184-3
43. Gerhart J, Danilchik M, Doniach T, Roberts S, Rowning B, et al. (1989) Cortical rotation of the *Xenopus* egg: Consequences for the anteroposterior pattern of embryonic dorsal development. *Development* 107 Suppl:37-51.
44. Ubbels GA, Hara K, Koster CH, Kirschner MW (1983) Evidence for a functional role of the cytoskeleton in determination of the dorsoventral axis in *Xenopus laevis* eggs. *J Embryol Exp Morph* 77:15-37.
45. Elinson RP, Rowning B (1988) A transient array of parallel microtubules in frog eggs: Potential tracks for a cytoplasmic rotation that specifies the dorso-ventral axis. *Dev Biol* 128:185-197. doi:10.1016/0012-1606(88)90281-3
46. Larabell CA, Rowning BA, Wells J, Wu M, Gerhart JC (1996) Confocal microscopy analysis of living *Xenopus* eggs and the mechanism of cortical rotation. *Development* 122:1281-1289.
47. Rowning BA, Wells J, Wu M, Gerhart JC, Moon RT, et al. (1997) Microtubule-mediated transport of organelles and localization of β -catenin to the future dorsal side of *Xenopus* eggs. *Proc Natl Acad Sci USA* 94:1224-1229. doi:10.1073/pnas.94.4.1224
48. Nusse R, Varmus H (2012) Three decades of Wnts: A personal perspective on how a scientific field developed. *EMBO J* 31:2670-2684. doi:10.1038/emboj.2012.146
49. Spemann H, Mangold H (1924) Über induktion von embryonalanlagen durch implantation artfremder organisatoren. *Arch Mikro Anat Entwicklungsmech* 100:599-638. doi:10.1007/BF02108133
50. Dumont JN (1972) Oogenesis in *Xenopus laevis* (Daudin): I. Stages of oocyte development in laboratory maintained animals. *J Morphol* 136(2):153-179. doi:10.1002/jmor.1051360203
51. Danilchik MV, Gerhart JC (1987) Differentiation of the animal-vegetal axis in *Xenopus laevis* oocytes: I. Polarized intracellular translocation of platelets establishes the yolk gradient. *Dev Biol* 122(1):101-112. doi:10.1016/0012-1606(87)90336-8
52. King ML, Messitt TJ, Mowry KL (2005) Putting RNAs in the right place at the right time: RNA localization in the frog oocyte. *Biol Cell* 97:19-33. doi:10.1042/BC20040067
53. Thomsen GH, Melton DA (1993) Processed Vg1 protein is an axial mesoderm inducer in *Xenopus*. *Cell* 74:433-441.
54. Mosquera L, Forristall C, Zhou Y, King ML (1993) A mRNA localized to the vegetal cortex of *Xenopus* oocytes encodes a protein with a nanos-like zinc finger domain. *Development* 117:377-386.
55. Yisraeli JK, Sokol S, Melton DA (1990) A two-step model for the localization of maternal mRNA in *Xenopus* oocytes: Involvement of microtubules and microfilaments in the translocation and anchoring of Vg1 mRNA. *Development* 108:289-298.
56. Forristall C, Pondel M, Chen L, King ML (1995) Patterns of localization and cytoskeletal association of two vegetally localized RNAs, *Vg1* and *Xcat-2*. *Development* 121:201-208.
57. Nijjar S, Woodland HR (2013) Localisation of RNAs into the germ plasm of vitellogenic *Xenopus* oocytes. *PLoS One* 8:e61847. doi:10.1371/journal.pone.0061847
58. Guraya SS (1979) Recent advances in the morphology, cytochemistry, and function of Balbiani's vitelline body in animal oocytes. *Int Rev Cytol* 59:249-321. doi:10.1016/S0074-7696(08)61664-2
59. al-Mukhtar KA, Webb AC (1971) An ultrastructural study of primordial germ cells, oögonia and early oocytes in *Xenopus laevis*. *J Embryol Exp Morphol* 26:195-217.

60. Heasman J, Quarmby J, Wylie CC (1984) The mitochondrial cloud of *Xenopus* oocytes: The source of germinal granule material. *Dev Biol* 105:458-489. doi:10.1016/0012-1606(84)90303-8
61. Cox RT, Spradling AC (2003) A Balbiani body and the fusome mediate mitochondrial inheritance during *Drosophila* oogenesis. *Development* 130:1579-1590. doi:10.1242/dev.00365
62. Chang P, Torres J, Lewis RA, Mowry KL, Houliston E, et al. (2004) Localization of RNAs to the mitochondrial cloud in *Xenopus* oocytes through entrapment and association with endoplasmic reticulum. *Mol Biol Cell* 15:4669-4681. doi:10.1091/mbc.E04-03-0265
63. Wilk K, Bilinski S, Dougherty MT, Kloc M (2004) Delivery of germinal granules and localized RNAs via the messenger transport organizer pathway to the vegetal cortex of *Xenopus* oocytes occurs through directional expansion of the mitochondrial cloud. *Int J Dev Biol* 49:17-21. doi:10.1387/ijdb.041906kw
64. Mowry KL, Melton DA (1992) Vegetal messenger RNA localization directed by a 340-nt RNA sequence element in *Xenopus* oocytes. *Science* 255:991-994. doi:10.1126/science.1546297
65. Bubunenko M, Kress TL, Vempati UD, Mowry KL, King ML (2002) A consensus RNA signal that directs germ layer determinants to the vegetal cortex of *Xenopus* oocytes. *Dev Biol* 248:82-92. doi:10.1006/dbio.2002.0719
66. Jansen R (2001) mRNA localization: Message on the move. *Nat Rev Mol Cell Biol* 2:247-256. doi:10.1038/35067016
67. Elinson RP, King ML, Forristall C (1993) Isolated vegetal cortex from *Xenopus* oocytes selectively retains localized mRNAs. *Dev Biol* 160:554-562. doi:10.1006/dbio.1993.1329
68. Kageura H (1997) Activation of dorsal development by contact between the cortical dorsal determinant and the equatorial core cytoplasm in eggs of *Xenopus laevis*. *Development* 124:1543-1551.
69. Alarcón VB, Elinson RP (2001) RNA anchoring in the vegetal cortex of the *Xenopus* oocyte. *J Cell Sci* 114:1731-1741.
70. Boveri T (1901) Die polarität von ovocyte, ei und larve des *Strongylocentrotus lividus*. *Zool Jahrb Abt Anat Ontog Tier* 14:630-653.
71. Kominami T (1983) Establishment of embryonic axes in larvae of the starfish, *Asterina pectinifera*. *J Embryol Exp Morphol* 75:87-100.
72. Schulze FE (1893) Über die bezeichnung der lage und richtung im thierkörper. *Biol Centralblatt* 13:1-7.
73. Hörstadius S (1973) *Experimental Embryology of Echinoderms*. Clarendon Press (Oxford).
74. Kominami T (1988) Determination of dorso-ventral axis in early embryos of the sea urchin, *Hemicentrotus pulcherrimus*. *Dev Biol* 127:187-196. doi:10.1016/0012-1606(88)90200-X
75. Cameron RA, Fraser SE, Britten RJ, Davidson EH (1989) The oral-aboral axis of a sea urchin embryo is specified by first cleavage. *Development* 106:641-647.
76. Henry JJ, Klueg KM, Raff RA (1992) Evolutionary dissociation between cleavage, cell lineage and embryonic axes in sea urchin embryos. *Development* 114:931-938.
77. Wikramanayake AH, Klein WH (1997) Multiple signaling events specify ectoderm and pattern the oral-aboral axis in the sea urchin embryo. *Development* 124:13-20.
78. Goldstein B, Freeman G (1997) Axis specification in development. *BioEssays* 19:105-116. doi:10.1002/bies.950190205
79. Coffman JA, McCarthy JJ, Dickey-Sims C, Robertson AJ (2004) Oral-aboral axis specification in the sea urchin embryo: II. Mitochondrial distribution and redox state contribute to establishing polarity in *Strongylocentrotus purpuratus*. *Dev Biol* 273:160-171. doi:10.1016/j.ydbio.2004.06.005
80. Flowers VL, Courteau GR, Poustka AJ, Weng W, Venuti JM (2004) Nodal/activin signaling establishes oral-aboral polarity in the early sea urchin embryo. *Dev Dyn* 231:727-740. doi:10.1002/dvdy.20194
81. Duboc V, Röttinger E, Besnardeau L, Lepage T (2004) Nodal and BMP2/4 signaling organizes the oral-aboral axis of the sea urchin embryo. *Dev Cell* 6:397-410. doi:10.1016/S1534-5807(04)00056-5
82. Bergeron K-F, Xu X, Brandhorst BP (2011) Oral-aboral patterning and gastrulation of sea urchin embryos depend on sulfated glycosaminoglycans. *Mech Dev* 128:71-89. doi:10.1016/j.mod.2010.11.001
83. Peterson KJ, Arenas-Mena C, Davidson EH (2000) The A/P axis in echinoderm ontogeny and evolution: Evidence from fossils and molecules. *Evol Dev* 2:93-101. doi:10.1046/j.1525-142x.2000.00042.x
84. Morris VB (2007) Origins of radial symmetry identified in an echinoderm during adult development and the inferred axes of ancestral bilateral symmetry. *Proc R Soc Lond B* 274:1511-1516. doi:10.1098/rspb.2007.0312
85. Maruyama YK, Nakaseko Y, Yagi S (1985) Localization of cytoplasmic determinants responsible for primary mesenchyme formation and gastrulation in the unfertilized eggs of the sea urchin *Hemicentrotus pulcherrimus*. *J Exp Zool* 236:155-163. doi:10.1002/jez.1402360206
86. Wikramanayake AH, Huang L, Klein WH (1998) β -Catenin is essential for patterning the maternally specified animal-vegetal axis in the sea urchin embryo. *Proc Natl Acad Sci USA* 95:9343-9348.
87. Logan CY, Miller JR, Ferkowicz MJ, McClay DR (1999) Nuclear β -catenin is required to specify vegetal cell fates in the sea urchin embryo. *Development* 126:345-357.
88. Angerer LM, Angerer RC (2000) Animal-vegetal axis patterning mechanisms in the early sea urchin embryo. *Dev Biol* 218:1-12. doi:10.1006/dbio.1999.9553
89. Brandhorst BP, Klein WH (2002) Molecular patterning along the sea urchin animal-vegetal axis. *Int Rev Cytol* 213:183-232. doi:10.1016/S0074-7696(02)13015-4
90. Di Carlo M, Romancino DP, Montana G, Ghersi G (1994) Spatial distribution of two maternal messengers in *Paracentrotus lividus* during oogenesis and embryogenesis. *Proc Natl Acad Sci USA* 91:5622-5626. doi:10.1073/pnas.91.12.5622
91. Vlahou A, Gonzalez-Rimbau M, Flytzanis CN (1996) Maternal mRNA encoding the orphan steroid receptor SpCOUP-TF is localized in sea urchin eggs. *Development* 122:521-526.
92. Schroeder TE (1980) The jelly canal marker of polarity for sea urchin oocytes, eggs, and embryos. *Exp Cell Res* 128:490-494. doi:10.1016/0014-4827(80)90088-9
93. Schroeder TE (1980) Expressions of the prefertilization polar axis in sea urchin eggs. *Dev Biol* 79:428-443. doi:10.1016/0012-1606(80)90128-1
94. Schroeder TE (1985) Cortical expressions of polarity in the starfish oocyte. *Dev Growth Differ* 27:311-321. doi:10.1111/j.1440-169X.1985.00311.x
95. Schroeder TE (1986) The egg cortex in early development of sea urchins and starfish. In: Browder LW, ed. *Developmental Biology*. Plenum Press (New York) v2, pp 59-100.
96. Otto JJ, Schroeder TE (1984) Microtubule arrays in the cortex and near the germinal vesicle of immature starfish oocytes. *Dev Biol* 101:274-281. doi:10.1016/0012-1606(84)90141-6
97. Bonder EM, Fishkind DJ, Cotran NM, Begg DA (1989) The cortical actin-membrane cytoskeleton of unfertilized sea urchin eggs: Analysis of the spatial organization and relationship of filamentous actin, nonfilamentous actin, and egg spectrin. *Dev Biol* 134:327-341. doi:10.1016/0012-1606(89)90105-X
98. Boyle JA, Ernst SG (1989) Sea urchin oocytes possess elaborate cortical arrays of microfilaments, microtubules, and intermediate filaments. *Dev Biol* 134:72-84. doi:10.1016/0012-1606(89)90079-1
99. Gorter E, Grendel F (1925) On bimolecular layers of lipoids on the chromocytes of the blood. *J Exp Med* 41:439-444. doi:10.1084/jem.41.4.439

100. Frye LD, Edidin M (1970) The rapid mixing of cell surface antigens after the formation of mouse-human heterokaryons. *J Cell Sci* 7:319-335.
101. Singer SJ, Nicolson GL (1972) The fluid mosaic model of the structure of cell membranes. *Science* 175:720-731. doi:10.1126/science.175.4023.720
102. Peters R, Peters J, Tews KH, Bähr W (1974) A micro-fluorimetric study of translational diffusion in erythrocyte membranes. *Biochim Biophys Acta* 367:282-294. doi:10.1016/0005-2736(74)90085-6
103. Edidin M, Zagyansky Y, Lardner TJ (1976) Measurement of membrane protein lateral diffusion in single cells. *Science* 191:466-468. doi:10.1126/science.1246629
104. Axelrod D, Ravdin P, Koppel DE, Schlessinger J, Webb WW, *et al.* (1976) Lateral motion of fluorescently labeled acetylcholine receptors in membranes of developing muscle fibers. *Proc Natl Acad Sci USA* 73:4594-4598. doi:10.1073/pnas.73.12.4594
105. Cherry RJ (1979) Rotational and lateral diffusion of membrane proteins. *Biochim Biophys Acta* 559(4):289-327. doi:10.1016/0304-4157(79)90009-1
106. Almers W, Stirling C (1984) Distribution of transport proteins over animal cell membranes. *J Membrane Biol* 77:169-186. doi:10.1007/BF01870567
107. Edidin M, Kuo SC, Sheetz MP (1991) Lateral movements of membrane glycoproteins restricted by dynamic cytoplasmic barriers. *Science* 254:1379-1382. doi:10.1126/science.1835798
108. Engleman DM (2005) Membranes are more mosaic than fluid. *Nature* 438:578-580. doi:10.1038/nature04394
109. Kusumi A, Nakada C, Ritchie K, Murase K, Suzuki K, *et al.* (2005) Paradigm shift of the plasma membrane concept from the two-dimensional continuum fluid to the partitioned fluid: High-speed single-molecule tracking of membrane molecules. *Annu Rev Biophys Biomol Struct* 34:351-378. doi:10.1146/annurev.biophys.34.040204.144637
110. Stier A, Sackmann E (1973) Spin labels as enzyme substrates: Heterogeneous lipid distribution in liver microsomal membranes. *Biochim Biophys Acta* 311:400-408. doi:10.1016/0005-2736(73)90320-9
111. Karnovsky MJ, Kleinfeld AM, Hoover RL, Klausner RD (1982) The concept of lipid domains in membranes. *J Cell Biol* 94:1-6. doi:10.1083/jcb.94.1.1
112. Simons K, van Meer G (1988) Lipid sorting in epithelial cells. *Biochem* 27:6197-6202. doi:10.1021/bi00417a001
113. Simons K, Ikonen E (1997) Functional rafts in cell membranes. *Nature* 387:569-572. doi:10.1038/42408
114. Braccia A, Villani M, Immerdal L, Niels-Christiansen LL, Nystrom BT, *et al.* (2003) Microvillar membrane microdomains exist at physiological temperature: Role of galectin-4 as lipid raft stabilizer revealed by "superrafts." *J Biol Chem* 278:15679-15684. doi:10.1074/jbc.M211228200
115. Douglass AD, Vale RD (2005) Single-molecule microscopy reveals plasma membrane microdomains created by protein-protein networks that exclude or trap signaling molecules in T cells. *Cell* 121:937-950. doi:10.1016/j.cell.2005.04.009
116. Nichols B (2005) Without a raft. *Nature* 436:638-639. doi:10.1038/436638a
117. Hancock JF (2006) Lipid rafts: Contentious only from simplistic standpoints. *Nat Rev Mol Cell Biol* 7:456-462. doi:10.1038/nrm1925
118. Eggeling C, Ringemann C, Medda R, Schwarzmann G, Sandhoff K, *et al.* (2009) Direct observation of the nanoscale dynamics of membrane lipids in a living cell. *Nature* 457:1159-1162. doi:10.1038/nature07596
119. He H-T, Marguet D (2011) Detecting nanodomains in living cell membrane by fluorescence correlation spectroscopy. *Annu Rev Phys Chem* 62:417-436. doi:10.1146/annurev-physchem-032210-103402
120. Loura LM, Prieto M (2011) FRET in membrane biophysics: An overview. *Front Physiol* 2:82. doi:10.3389/fphys.2011.00082
121. Owen DM, Magenau A, Williamson D, Gaus K (2012) The lipid raft hypothesis revisited—new insights on raft composition and function from super-resolution fluorescence microscopy. *BioEssays* 34:739-747. doi:10.1002/bies.201200044
122. Sonnino S, Prinetti A (2013) Membrane domains and the "lipid raft" concept. *Curr Med Chem* 20:4-21. doi:10.2174/0929867311320010003
123. Chichili GR, Rodgers W (2009) Cytoskeleton-membrane interactions in membrane raft structure. *Cell Mol Life Sci* 66:2319-2328. doi:10.1007/s00018-009-0022-6
124. Head BP, Patel HH, Insel PA (2013) Interaction of membrane/lipid rafts with the cytoskeleton: Impact on signaling and function. *Biochim Biophys Acta pii:S0005-2736(13)00258-7*. doi:10.1016/j.bbamem.2013.07.018
125. Simons K, Sampaio JL (2011) Membrane organization and lipid rafts. *Cold Spring Harb Perspect Biol* 3(10):a004697. doi:10.1101/cshperspect.a004697
126. Reeves VL, Thomas CM, Smart EJ (2012) Lipid rafts, caveolae and GPI-linked proteins. *Adv Exp Med Biol* 729:3-13. doi:10.1007/978-1-4614-1222-9_1
127. Suzuki KGN (2012) Lipid rafts generate digital-like signal transduction in cell plasma membranes. *Biotechnol J* 7:753-761. doi:10.1002/biot.201100360
128. Bashirullah A, Cooperstock RL, Lipshitz HD (1998) RNA localization in development. *Annu Rev Biochem* 67:335-394. doi:10.1146/annurev.biochem.67.1.335
129. Palacios IM (2007) How does an mRNA find its way? Intracellular localisation of transcripts. *Semin Cell Dev Biol* 18:163-170. doi:10.1016/j.semcdb.2007.01.008
130. Lécuyer E, Yoshida H, Parthasarathy N, Alm C, Babak T, *et al.* (2007) Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and function. *Cell* 131:174-187. doi:10.1016/j.cell.2007.08.003
131. Holt CE, Bullock SL (2009) Subcellular mRNA localization in animal cells and why it matters. *Science* 326:1212-1216. doi:10.1126/science.1176488
132. Singer RH (1993) RNA zipcodes for cytoplasmic addresses. *Curr Biol* 3:719-721. doi:10.1016/0960-9822(93)90079-4
133. Wilhelm JE, Vale RD (1993) RNA on the move: The mRNA localization pathway. *J Cell Biol* 123:269-274. doi:10.1083/jcb.123.2.269
134. Lawrence JB, Singer RH (1986) Intracellular localization of messenger RNAs for cytoskeletal proteins. *Cell* 45:407-415. doi:10.1016/0092-8674(86)90326-0
135. Ross AF, Oleynikov Y, Kislauskis EH, Raneja KL, Singer RH (1997) Characterization of a β -actin mRNA zipcode-binding protein. *Mol Cell Biol* 17:2158-2165.
136. Katz ZB, Wells AL, Park HY, Wu B, Shenoy SM, *et al.* (2012) β -actin mRNA compartmentalization enhances focal adhesion stability and directs cell migration. *Genes Dev* 26:1885-1890. doi:10.1101/gad.190413.112
137. Havin L, Git A, Elisha Z, Oberman F, Yaniv K, *et al.* (1998) RNA-binding protein conserved in both microtubule- and microfilament-based RNA localization. *Genes Dev* 12:1593-1598. doi:10.1101/gad.12.11.1593
138. Muslimov IA, Patel MV, Rose A, Tiedge H (2011) Spatial code recognition in neuronal RNA targeting: Role of RNA-hnRNP A2 interactions. *J Cell Biol* 194:441-457. doi:10.1083/jcb.201010027
139. Williams LT (1989) Signal transduction by the platelet-derived growth factor receptor. *Science* 243:1564-1570. doi:10.1126/science.2538922
140. Latham VM Jr, Kislauskis EH, Singer RH, Ross AF (1994) β -actin mRNA localization is regulated by signal transduction mechanisms. *J Cell Biol* 126:1211-1219. doi:10.1083/jcb.126.5.1211

141. Hubbard SR, Till JH (2000) Protein tyrosine kinase structure and function. *Annu Rev Biochem* 69:373-398. doi:10.1146/annurev.biochem.69.1.373
142. Lemmon MA, Schlessinger J (2010) Cell signaling by receptor tyrosine kinases. *Cell* 141:1117-1134. doi:10.1016/j.cell.2010.06.011
143. Wittmann T, Waterman-Storer CM (2001) Cell motility: Can Rho GTPases and microtubules point the way? *J Cell Sci* 114:3795-3803.
144. Latham VM, Yu EH, Tullio AN, Adelstein RS, Singer RH (2001) A Rho-dependent signaling pathway operating through myosin localizes beta-actin mRNA in fibroblasts. *Curr Biol* 11:1010-1016. doi:10.1016/S0960-9822(01)00291-3
145. Etienne-Manneville S, Hall A (2003) Cdc42 regulates GSK-3beta and adenomatous polyposis coli to control cell polarity. *Nature* 421:753-756. doi:10.1038/nature01423
146. Iden S, Collard JG (2008) Crosstalk between small GTPases and polarity proteins in cell polarization. *Nat Rev Mol Cell Biol* 9:846-859. doi:10.1038/nrm2521
147. Martin-Belmonte F, Mostov K (2008) Regulation of cell polarity during epithelial morphogenesis. *Curr Opin Cell Biol* 20:227-234. doi:10.1016/j.ceb.2008.01.001
148. Apodaca G, Gallo LI, Bryant DM (2012) Role of membrane traffic in the generation of epithelial cell asymmetry. *Nat Cell Biol* 14:1235-1243. doi:10.1038/ncb2635
149. Hay ED (2005) The mesenchymal cell, its role in the embryo, and the remarkable signaling mechanisms that create it. *Dev Dyn* 233:706-720. doi:10.1002/dvdy.20345
150. Bryant DM, Mostov KE (2008) From cells to organs: Building polarized tissue. *Nat Rev Mol Cell Biol* 9:887-901. doi:10.1038/nrm2523
151. Huang X, Saint-Jeannet J-P (2004) Induction of the neural crest and the opportunities of life on the edge. *Dev Biol* 275:1-11. doi:10.1016/j.ydbio.2004.07.033
152. Kalluri R, Weinberg RA (2009) The basics of epithelial-mesenchymal transition. *J Clin Invest* 119:1420-1428. doi:10.1172/JCI39104
153. Kerosuo L, Bronner-Fraser M (2012) What is bad in cancer is good in the embryo: Importance of EMT in neural crest development. *Semin Cell Dev Biol* 23:320-332. doi:10.1016/j.semcdb.2012.03.010
154. Yeaman C, Grindstaff KK, Nelson WJ (1999) New perspectives on mechanisms involved in generating epithelial cell polarity. *Physiol Rev* 79:73-98.
155. Nejsum LN, Nelson WJ (2009) Epithelial cell surface polarity: The early steps. *Front Biosci* 14:1088-1098. doi:10.2741/3295
156. Gassama-Diagne A, Yu W, ter Beest M, Martin-Belmonte F, Kierbel A, et al. (2006) Phosphatidylinositol-3,4,5-trisphosphate regulates the formation of the basolateral plasma membrane in epithelial cells. *Nat Cell Biol* 8:963-970. doi:10.1038/ncb1461
157. Martin-Belmonte F, Gassama A, Datta A, Yu W, Rescher U, et al. (2007) PTEN-mediated apical segregation of phosphoinositides controls epithelial morphogenesis through Cdc42. *Cell* 128:383-397. doi:10.1016/j.cell.2006.11.051
158. Brown DA, Rose JK (1992) Sorting of GPI-anchored proteins to glycolipid-enriched membrane subdomains during transport to the apical cell surface. *Cell* 68:533-544.
159. Cao X, Surma MA, Simons K (2012) Polarized sorting and trafficking in epithelial cells. *Cell Res* 22:793-805. doi:10.1038/cr.2012.64
160. Ridley AJ, Schwartz MA, Burridge K, Firtel RA, Ginsberg MH, et al. (2003) Cell migration: Integrating signals from front to back. *Science* 302:1704-1709. doi:10.1126/science.1092053
161. Nelson WJ (2009) Remodeling epithelial cell organization: Transitions between front-rear and apical-basal polarity. *Cold Spring Harb Perspect Biol* 1:a000513. doi:10.1101/cshperspect.a000513
162. Varki A (2007) Glycan-based interactions involving vertebrate sialic-acid-recognizing proteins. *Nature* 446:1023-1029. doi:10.1038/nature05816
163. Furukawa K, Ohkawa Y, Yamauchi Y, Hamamura K, Ohmi Y, et al. (2012) Fine tuning of cell signals by glycosylation. *J Biochem* 151:573-578. doi:10.1093/jb/mvs043
164. Schnaar RL (1985) The membrane is the message. *The Sciences* (May-June):34-40.
165. Laine RA (1994) A calculation of all possible oligosaccharide isomers both branched and linear yields 1.05×10^{12} structures for a reducing hexasaccharide. *Glycobiology* 4:759-767. doi:10.1093/glycob/4.6.759
166. Gabius H-J (2000) Biological information transfer beyond the genetic code: The sugar code. *Naturwissenschaften* 87:108-121. doi:10.1007/s001140050687
167. Gabius H-J, Siebert H-C, André S, Jiménez-Barbero J, Rüdiger H (2004) Chemical biology of the sugar code. *Chembiochem* 5:740-764. doi:10.1002/cbic.200300753
168. Cummings RD (2009) The repertoire of glycan determinants in the human glycome. *Mol Biosyst* 5:1077-1248. doi:10.1039/b907931a
169. Rüdiger H, Gabius H-J (2009) The biochemical basis and coding capacity of the sugar code. In: Gabius H-J, ed. *The Sugar Code: Fundamentals of Glycosciences*. Wiley-VCH Verlag (Weinheim, Germany) pp 3-13.
170. Gabius H-J (2006) Cell surface glycans: The why and how of their functionality as biochemical signals in lectin-mediated information transfer. *Crit Rev Immunol* 26:43-79. doi:10.1615/CritRevImmunol.v26.i1.30
171. Nugent MA (2000) Heparin sequencing brings structure to the function of complex oligosaccharides. *Proc Natl Acad Sci USA* 97:10301-10303. doi:10.1073/pnas.97.19.10301
172. Buddecke E (2009) Proteoglycans. In: Gabius H-J, ed. *The Sugar Code: Fundamentals of Glycosciences*. Wiley-VCH Verlag (Weinheim, Germany) pp 199-216.
173. Turnbull JE (2010) Heparan sulfate glycomics: Towards systems biology strategies. *Biochem Soc Trans* 38:1356-1360. doi:10.1042/BST0381356
174. Feizi T (1985) Demonstration by monoclonal antibodies that carbohydrate structures of glycoproteins and glycolipids are onco-developmental antigens. *Nature* 314:53-57. doi:10.1038/314053a0
175. Rastan S, Thorpe SJ, Scudder P, Brown S, Gooi HC, et al. (1985) Cell interactions in preimplantation embryos: Evidence for involvement of saccharides of the poly-N-acetyllactosamine series. *J Embryol Exp Morph* 87:115-128.
176. Pennington JE, Rastan S, Roelke D, Feizi T (1985) Saccharide structures of the mouse embryo during the first eight days of development. *J Embryol Exp Morph* 90:335-361.
177. Fenderson BA, Eddy EM, Hakomori S (1990) Glycoconjugate expression during embryogenesis and its biological significance. *BioEssays* 12:173-179. doi:10.1002/bies.950120406
178. Thorpe S, Bellairs R, Feizi T (1988) Developmental patterning of carbohydrate antigens during early embryogenesis of the chick: Expression of antigens of the poly-N-acetyllactosamine series. *Development* 102:193-210.
179. Loveless W, Bellairs R, Thorpe SJ, Page M, Feizi T (1990) Developmental patterning of the carbohydrate antigen FC10.2 during early embryogenesis in the chick. *Development* 108:97-106.
180. Blum AS, Barnstable CJ (1987) O-Acetylation of a cell-surface carbohydrate creates discrete molecular patterns during neural development. *Proc Natl Acad Sci USA* 84:8716-8720. doi:10.1073/pnas.84.23.8716
181. Perrimon N, Bernfield M (2000) Specificities of heparan sulphate proteoglycans in developmental processes. *Nature* 404:725-728. doi:10.1038/35008000
182. Haltiwanger RS, Lowe JB (2004) Role of glycosylation in development. *Annu Rev Biochem* 73:491-537. doi:10.1146/annurev.biochem.73.011303.074043

183. Topczewski J, Sepich DS, Myers DC, Walker C, Amores A, *et al.* (2001) The zebrafish glypican knypek controls cell polarity during gastrulation movements of convergent extension. *Dev Cell* 1:251-64. doi:10.1016/S1534-5807(01)00005-3
184. Turnbull J, Powell A, Guimond S (2001) Heparan sulfate: Decoding a dynamic multifunctional cell regulator. *Trends Cell Biol* 11:75-82. doi:10.1016/S0962-8924(00)01897-3
185. Princivalle M, de Agostini A (2002) Developmental roles of heparan sulfate proteoglycans: A comparative review in *Drosophila*, mouse and human. *Int J Dev Biol* 46:267-278.
186. Bülow HE, Hobert O (2006) The molecular diversity of glycosaminoglycans shapes animal development. *Annu Rev Cell Dev Biol* 22:375-407. doi:10.1146/annurev.cellbio.22.010605.093433
187. Hillenkamp F, Karas M, Beavis RC, Chait BT (1991) Matrix-assisted laser desorption/ionization mass spectrometry of biopolymers. *Anal Chem* 63:1193A-1203A. doi:10.1021/ac00024a002
188. Lopez PHH, Schnaar RL (2009) Gangliosides in cell recognition and membrane protein regulation. *Curr Opin Struct Biol* 19:549-557. doi:10.1016/j.sbi.2009.06.001
189. Sharon N, Lis H (2004) History of lectins: From hemagglutinins to biological recognition molecules. *Glycobiology* 14:53R-62R. doi:10.1093/glycob/cwh122
190. Drickamer K, Taylor ME (1993) Biology of animal lectins. *Annu Rev Cell Biol* 9:237-264. doi:10.1146/annurev.cb.09.110193.001321
191. Gabius H-J (1997) Animal lectins. *Eur J Biochem* 243:543-576. doi:10.1111/j.1432-1033.1997.t01-1-00543.x
192. Brandley BK, Schnaar RL (1986) Cell-surface carbohydrates in cell recognition and response. *J Leukocyte Biol* 40:97-111.
193. Fukuda M, Hiraoka N, Yeh J-C (1999) C-type lectins and sialyl Lewis^x oligosaccharides: Versatile roles in cell-cell interaction. *J Cell Biol* 147:467-470. doi:10.1083/jcb.147.3.467
194. Gabius H-J (2009) Animal and human lectins. In: Gabius H-J, ed. *The Sugar Code: Fundamentals of Glycosciences*. Wiley-VCH Verlag (Weinheim) pp 317-328.
195. Griffith CM, Sanders EJ (1991) Changes in glycoconjugate expression during early chick embryo development: A lectin-binding study. *Anat Rec* 231:238-250. doi:10.1002/ar.1092310212
196. Lipscomb BW, Treloar HB, Klenoff J, Greer CA (2003) Cell surface carbohydrates and glomerular targeting of olfactory sensory neuron axons in the mouse. *J Comp Neurol* 467:22-31. doi:10.1002/cne.10910
197. Barondes SH, Castronovo V, Cooper DN, Cummings RD, Drickamer K, *et al.* (1994) Galectins: A family of animal β -galactoside-binding lectins. *Cell* 76:597-598. doi:10.1016/0092-8674(94)90498-7
198. Kasai K, Hirabayashi J (1996) Galectins: A family of animal lectins that decipher glycodes. *J Biochem* 119:1-8. doi:10.1093/oxfordjournals.jbchem.a021192
199. Cooper DNW (2002) Galectinomics: Finding themes in complexity. *Biochim Biophys Acta* 1572:209-231. doi:10.1016/S0304-4165(02)00310-0
200. Van Den Brùle FA, Fernandez PL, Buicu C, Liu F-T, Jackers P, *et al.* (1997) Differential expression of galectin-1 and galectin-3 during first trimester human embryogenesis. *Dev Dyn* 209:399-405.
201. Shoji H, Nishi N, Hirashima M, Nakamura T (2003) Characterization of the *Xenopus* galectin family: Three structurally different types as in mammals and regulated expression during embryogenesis. *J Biol Chem* 278:12285-12293. doi:10.1074/jbc.M209008200
202. Rabinovich GA, Toscano MA, Jackson SS, Vasta GR (2007) Functions of cell surface galectin-glycoprotein lattices. *Curr Opin Struct Biol* 17:513-520. doi:10.1016/j.sbi.2007.09.002
203. Garner OB, Baum LG (2008) Galectin-glycan lattices regulate cell-surface glycoprotein organization and signaling. *Biochem Soc Trans* 36:1472-1477. doi:10.1042/BST0361472
204. Bhat R, Lerea KM, Peng H, Kaltner H, Gabius H-J, *et al.* (2011) A regulatory network of two galectins mediates the earliest steps of avian limb skeletal morphogenesis. *BMC Dev Biol* 11:6. doi:10.1186/1471-213X-11-6
205. Broom MF, Shepherd MG, Sullivan PA (1991) Changes in cell envelope glycoproteins during germ-tube formation of *Candida albicans*. *Microbios* 67(274):7-21.
206. Foster AJ, Bird RA, Kelly SL, Nishimura K, Poyner D, *et al.* (2004) FITC-lectin avidity of *Cryptococcus neoformans* cell wall and capsular components. *Mycologia* 96:1-8. doi:10.2307/3761980
207. Wanchoo A, Lewis MW, Keyhani NO (2009) Lectin mapping reveals stage-specific display of surface carbohydrates in *in vitro* and haemolymph-derived cells of the entomopathogenic fungus *Beauveria bassiana*. *Microbiology* 155:3121-3133. doi:10.1099/mic.0.029157-0
208. Cattaneo F, Pasini ME, Perotti ME (1997) Glycosidases are present on the surface of *Drosophila melanogaster* spermatozoa. *Mol Reprod Dev* 48:276-281.
209. Intra J, Cenni F, Perotti ME (2006) An alpha-L-fucosidase potentially involved in fertilization is present on *Drosophila* spermatozoa surface. *Mol Reprod Dev* 73:1149-1158. doi:10.1002/mrd.20425
210. Perotti ME, Riva A (1988) Concanavalin A binding sites on the surface of *Drosophila melanogaster* sperm: A fluorescence and ultrastructural study. *J Ultrastruct Mol Struct Res* 1988 Aug;100(2):173-182. doi:10.1016/0889-1605(88)90024-9
211. Perotti ME, Camplani A, Faré C (1990) The carbohydrates of the envelopes of *Drosophila* egg before and after fertilization. *J Cell Biol* 111(5):486a.
212. Fabian L, Brill JA (2012) *Drosophila* spermiogenesis: Big things come from little packages. *Spermatogenesis* 2:197-212. doi:10.4161/spmg.21798
213. Prescher JA, Dube DH, Bertozzi CR (2004) Chemical remodeling of cell surfaces in living animals. *Nature* 430:873-877. doi:10.1038/nature02791
214. Campbell CT, Sampathkumar S-G, Yarema KJ (2007) Metabolic oligosaccharide engineering: Perspectives, applications, and future directions. *Mol Biosyst* 3:187-194. doi:10.1039/b614939c
215. Bernardi A, Cheshev P (2008) Interfering with the sugar code: Design and synthesis of oligosaccharide mimics. *Chem Eur J* 14:7434-7441. doi:10.1002/chem.200800597
216. Laughlin ST, Baskin JM, Amacher SL, Bertozzi CR (2008) In vivo imaging of membrane-associated glycans in developing zebrafish. *Science* 320:664-667. doi:10.1126/science.1155106
217. Han E, Ding L, Ju H (2011) Highly sensitive fluorescent analysis of dynamic glycan expression on living cells using glyconanoparticles and functionalized quantum dots. *Anal Chem* 83:7006-7012. doi:10.1021/ac201488x
218. Cao J-T, Chen Z-X, Hao X-Y, Zhang P-H, Zhu J-J (2012) Quantum dots-based immunofluorescent microfluidic chip for the analysis of glycan expression at single cells. *Anal Chem* 84:10097-10104. doi:10.1021/ac302609y
219. Teorell T (1935) Studies on the "diffusion effect" upon ionic distribution. *Proc Natl Acad Sci USA* 21:152-161. doi:10.1073/pnas.21.3.152
220. Burr HS, Northrop FSC (1939) Evidence for the existence of an electro-dynamic field in living organisms. *Proc Natl Acad Sci USA* 25:284-288. doi:10.1073/pnas.25.6.284
221. Romanoff AL (1944) Bio-electric potentials and vital activity of the egg. *Biodynamica* 4:329-358.
222. Jaffe LF, Nuccitelli R (1977) Electrical controls of development. *Annu Rev Biophys Bioeng* 6:445-476. doi:10.1146/annurev.bb.06.060177.002305
223. Skou JC (1957) The influence of some cations on an adenosine triphosphatase from peripheral nerves. *Biochim Biophys Acta* 23:394-401. doi:10.1016/0006-3002(57)90343-8

224. Skou JC (1990) The energy coupled exchange of Na⁺ for K⁺ across the cell membrane. The Na⁺, K⁺-pump. *FEBS Lett* 268:314-324. doi:10.1016/0014-5793(90)81278-V
225. Hodgkin AL, Huxley AF (1952) A quantitative description of membrane current and its application to conduction and excitation in a nerve. *J Physiol* 117:500-544.
226. Ketchum KA, Joiner WJ, Sellers AJ, Kaczmarek LK, Goldstein SA (1995) A new family of outwardly rectifying potassium channel proteins with two pore domains in tandem. *Nature* 376:690-695. doi:10.1038/376690a0
227. Goldstein SA, Bockenhauer D, O'Kelly I, Zilberberg N (2001) Potassium leak channels and the KCNK family of two-P-domain subunits. *Nat Rev Neurosci* 2:175-184. doi:10.1038/35058574
228. McCaig CD, Rajnicsek AM, Song B, Zhao M (2005) Controlling cell behavior electrically: Current views and future potential. *Physiol Rev* 85:943-978. doi:10.1152/physrev.00020.2004
229. Nuccitelli R (2006) Endogenous electric fields in animals. In: *Handbook of Biological Effects of Electromagnetic Fields*. Third Edition. Taylor & Francis (Abingdon, U.K.) ch 2.
230. Barker AT, Jaffe LF, Vanable JW Jr (1982) The glabrous epidermis of cavies contains a powerful battery. *Am J Physiol* 242:R358-R366.
231. McCaig CD, Robinson KR (1982) The ontogeny of the trans-epidermal potential difference in frog embryos. *Dev Biol* 90:335-339. doi:10.1016/0012-1606(82)90382-7
232. Ling G, Gerard RW (1949) The normal membrane potential of frog sartorius fibers. *J Cell Comp Physiol* 34:383-396. doi:10.1002/jcp.1030340304
233. Neher E, Sakmann B (1976) Single-channel currents recorded from membrane of denervated frog muscle fibres. *Nature* 260:799-802. doi:10.1038/260799a0
234. Li WC, Soffe SR, Roberts A (2004) A direct comparison of whole cell patch and sharp electrodes by simultaneous recording from single spinal neurons in frog tadpoles. *J Neurophysiol* 92:380-386. doi:10.1152/jn.01238.2003
235. Förster T (1948) Zwischenmolekulare energiewandlung und fluoreszenz. *Annalen der Physik* 437:55-75. doi:10.1002/andp.19484370105
236. González JE, Oades K, Leychik Y, Harootunian A, Negulescu PA (1999) Cell-based assays and instrumentation for screening ion-channel targets. *Drug Discov Today* 4:431-439. doi:10.1016/S1359-6446(99)01383-5
237. Blackiston D, Adams DS, Lemire JM, Lobikin M, Levin M (2011) Transmembrane potential of GlyCl-expressing instructor cells induces a neoplastic-like conversion of melanocytes via a serotonergic pathway. *Dis Model Mech* 4:67-85. doi:10.1242/dmm.005561
238. Jaffe LF, Nuccitelli R (1974) An ultrasensitive vibrating probe for measuring steady extracellular currents. *J Cell Biol* 63:614-628. doi:10.1083/jcb.63.2.614
239. Hotary KB, Nuccitelli R, Robinson KR (1992) A computerized 2-dimensional vibrating probe for mapping extracellular current patterns. *J Neurosci Meth* 43:55-67. doi:10.1016/0165-0270(92)90067-N
240. Reid B, Nuccitelli R, Zhao M (2007) Non-invasive measurement of bioelectric currents with a vibrating probe. *Nat Protoc* 2:661-669. doi:10.1038/nprot.2007.91
241. Wallace RA, Steinhardt RA (1977) Maturation of *Xenopus* oocytes: II. Observations on membrane potential. *Dev Biol* 57:305-316. doi:10.1016/0012-1606(77)90217-2
242. Robinson KR (1979) Electrical currents through full-grown and maturing *Xenopus* oocytes. *Proc Natl Acad Sci USA* 76:837-841. doi:10.1073/pnas.76.2.837
243. Kline D, Robinson KR, Nuccitelli R (1983) Ion currents and membrane domains in the cleaving *Xenopus* egg. *J Cell Biol* 97:1753-1761. doi:10.1083/jcb.97.6.1753
244. Regen CM, Steinhardt RA (1986) Global properties of the *Xenopus* blastula are mediated by a high-resistance epithelial seal. *Dev Biol* 113:147-154. doi:10.1016/0012-1606(86)90117-X
245. Robinson KR, Stump RF (1984) Self-generated electrical currents through *Xenopus* neurulae. *J Physiol* 352:339-352.
246. Metcalf MEM, Shi R, Borgens RB (1994) Endogenous ionic currents and voltages in amphibian embryos. *J Exp Zool* 268:307-322. doi:10.1002/jez.1402680407
247. Hotary KB, Robinson KR (1991) The neural tube of the *Xenopus* embryo maintains a potential difference across itself. *Dev Brain Res* 59:65-73. doi:10.1016/0165-3806(91)90030-M
248. Jaffe LF (1981) The role of ionic currents in establishing developmental pattern. *Phil Trans R Soc Lond B* 295:553-556. doi:10.1098/rstb.1981.0160
249. Robinson KR (1983) Endogenous electrical current leaves the limb and pre-limb region of the *Xenopus* embryo. *Dev Biol* 97:203-211. doi:10.1016/0012-1606(83)90077-5
250. Jaffe LF, Stern CD (1979) Strong electrical currents leave the primitive streak of chick embryos. *Science* 206:569-571. doi:10.1126/science.573921
251. Kucera P, de Ribaupierre Y (1989) Extracellular electrical currents in the chick blastoderm. *Biol Bull Suppl* 176:118-122. doi:10.2307/1541659
252. Hotary KB, Robinson KR (1990) Endogenous electrical currents and the resultant voltage gradients in the chick embryo. *Dev Biol* 140:149-160. doi:10.1016/0012-1606(90)90062-N
253. Winkel GK, Richard Nuccitelli R (1989) Large ionic currents leave the primitive streak of the 7.5-day mouse embryo. *Biol Bull Suppl* 176:110-117. doi:10.2307/1541658
254. Altizer AM, Moriarty LJ, Bell SM, Schreiner CM, Scott WJ, et al. (2001) Endogenous electric current is associated with normal development of the vertebrate limb. *Dev Dyn* 221:391-401. doi:10.1002/dvdy.1158
255. Cooper MS, Schliwa M (1985) Electrical and ionic controls of tissue cell locomotion in DC electric fields. *J Neurosci Res* 13:223-244. doi:10.1002/jnr.490130116
256. Stump RF, Robinson KR (1983) *Xenopus* neural crest cell migration in an applied electrical field. *J Cell Biol* 97:1226-1233. doi:10.1083/jcb.97.4.1226
257. Robinson KR (1985) The responses of cells to electrical fields: A review. *J Cell Biol* 101:2023-2027. doi:10.1083/jcb.101.6.2023
258. Erickson CA, Nuccitelli R (1984) Embryonic fibroblast motility and orientation can be influenced by physiological electric fields. *J Cell Biol* 98:296-307. doi:10.1083/jcb.98.1.296
259. Nuccitelli R (1988) Physiological electric fields can influence cell motility, growth, and polarity. *Adv Cell Biol* 2:213-233. doi:10.1016/S1569-2558(08)60435-X
260. McKasson MJ, Huang L, Robinson KR (2008) Chick embryonic Schwann cells migrate anodally in small electrical fields. *Exp Neurol* 211:585-587. doi:10.1016/j.expneurol.2008.02.015
261. Stern CD (1981) Behaviour and motility of cultured chick mesoderm cells in steady electrical fields. *Exp Cell Res* 136:343-350. doi:10.1016/0014-4827(81)90013-6
262. Cormie P, Robinson KR (2007) Embryonic zebrafish neuronal growth is not affected by an applied electric field in vitro. *Neurosci Lett* 411:128-132. doi:10.1016/j.neulet.2006.10.030
263. Jaffe LF, Poo M-M (1979) Neurites grow faster towards the cathode than the anode in a steady field. *J Exp Zool* 209:115-128. doi:10.1002/jez.1402090114
264. Hinkle L, McCaig CD, Robinson KR (1981) The direction of growth of differentiating neurons and myoblasts from frog embryos in an applied electric field. *J Physiol* 314:121-135.
265. McCaig CD, Song B, Rajnicsek AM (2009) Electrical dimensions in cell science. *J Cell Sci* 122:4267-4276. doi:10.1242/jcs.023564
266. Hotary KB, Robinson KR (1992) Evidence of a role for endogenous electrical fields in chick embryo development. *Development* 114:985-996.

267. Hotary KB, Robinson KR (1994) Endogenous electrical currents and voltage gradients in *Xenopus* embryos and the consequences of their disruption. *Dev Biol* 166:789-800. doi:10.1006/dbio.1994.1357
268. Shi R, Borgens RB (1994) Embryonic neuroepithelial sodium transport, the resulting physiological potential, and cranial development. *Dev Biol* 165:105-116. doi:10.1006/dbio.1994.1238
269. Borgens RB, Shi R (1995) Uncoupling histogenesis from morphogenesis in the vertebrate embryo by collapse of the transneural tube potential. *Dev Dyn* 203:456-467. doi:10.1002/aja.1002030408
270. Harvey WR, Wiczorek H (1997) Animal plasma membrane energization by chemiosmotic H⁺ V-ATPases. *J Exp Biol* 200:203-216.
271. Nelson N, Perzov N, Cohen A, Hagai K, Padler V, *et al.* (2000) The cellular biology of proton-motive force generation by V-ATPases. *J Exp Biol* 203:89-95.
272. Hinton A, Bond S, Forgac M (2009) V-ATPase functions in normal and disease processes. *Pflugers Arch* 457:589-598. doi:10.1007/s00424-007-0382-4
273. Adams DS, Robinson KR, Fukumoto T, Yuan S, Albertson RC, *et al.* (2006) Early, H⁺-V-ATPase-dependent proton flux is necessary for consistent left-right patterning of non-mammalian vertebrates. *Development* 133:1657-1671. doi:10.1242/dev.02341
274. Adams DS, Masi A, Levin M (2007) H⁺ pump-dependent changes in membrane voltage are an early mechanism necessary and sufficient to induce *Xenopus* tail regeneration. *Development* 134:1323-1335. doi:10.1242/dev.02812
275. Cruciat CM, Ohkawara B, Acebron SP, Karaulanov E, Reinhard C, *et al.* (2010) Requirement of prorenin receptor and vacuolar H⁺-ATPase-mediated acidification for Wnt signaling. *Science* 327:459-463. doi:10.1126/science.1179802
276. Finbow ME, Harrison M, Jones P (1995) Ductin—a proton pump component, a gap junction channel and a neurotransmitter release channel. *BioEssays* 17:247-255. doi:10.1002/bies.950170311
277. Vandenberg LN, Morrie RD, Adams DS (2011) V-ATPase-dependent ectodermal voltage and pH regionalization are required for craniofacial morphogenesis. *Dev Dyn* 240:1889-1904. doi:10.1002/dvdy.22685
278. Beg AA, Jorgensen EM (2003) EXP-1 is an excitatory GABA-gated cation channel. *Nat Neurosci* 6:1145-1152. doi:10.1038/nn1136
279. Davies DL, Trudell JR, Mihic SJ, Crawford DK, Alkana RL (2003) Ethanol potentiation of glycine receptors expressed in *Xenopus* oocytes antagonized by increased atmospheric pressure. *Alcohol Clin Exp Res* 27:743-755. doi:10.1097/01.ALC.0000065722.31109.A1
280. Shan Q, Haddrill JL, Lynch JW (2001) Ivermectin, an unconventional agonist of the glycine receptor chloride channel. *J Biol Chem* 276:12556-12564. doi:10.1074/jbc.M011264200
281. Pai VP, Aw S, Shomrat T, Lemire JM, Levin M (2012) Transmembrane voltage potential controls embryonic eye patterning in *Xenopus laevis*. *Development* 139:313-323. doi:10.1242/dev.073759
282. Nuccitelli R (1988) Ionic currents in morphogenesis. *Experientia* 44:657-666. doi:10.1007/BF01941026
283. Shi R, Borgens RB (1995) Three-dimensional gradients of voltage during development of the nervous system as invisible coordinates for the establishment of embryonic pattern. *Dev Dyn* 202:101-114. doi:10.1002/aja.1002020202
284. Gurwitsch AG (1922) Über den begriff des embryonalen feldes. *Arch Entwicklung Org* 51:388-415. doi:10.1007/BF02554452
285. Davidson EH (1993) Later embryogenesis: Regulatory circuitry in morphogenetic fields. *Development* 118:665-690.
286. Belousov LV, Opitz JM, Gilbert SF (1997) Life of Alexander G. Gurwitsch and his relevant contribution to the theory of morphogenetic fields. *Int J Dev Biol* 41:771-779.
287. Levin M (2012) Morphogenetic fields in embryogenesis, regeneration, and cancer: Non-local control of complex patterning. *BioSystems* 109:243-261. doi:10.1016/j.biosystems.2012.04.005
288. Tseng A, Levin M (2013) Cracking the bioelectric code: Probing endogenous ionic controls of pattern formation. *Commun Integr Biol* 6(1):e22595. doi:10.4161/cib.22595
289. Wang ET, Sandberg R, Luo S, Khrebukova I, Zhang L, *et al.* (2008) Alternative isoform regulation in human tissue transcriptomes. *Nature* 456:470-476. doi:10.1038/nature07509
290. Pan Q, Shai O, Lee LJ, Frey BJ, Blencowe BJ (2008) Deep surveying of alternative splicing complexity in the human transcriptome by high-throughput sequencing. *Nat Genet* 40:1413-1415. doi:10.1038/ng.259
291. Barash Y, Calarco JA, Gao W, Pan Q, Wang X, *et al.* (2010) Deciphering the splicing code. *Nature* 465:53-59. doi:10.1038/nature09000
292. Kornblihtt AR, Schor IE, Alló M, Dujardin G, Petrillo E, *et al.* (2013) Alternative splicing: A pivotal step between eukaryotic transcription and translation. *Nat Rev Mol Cell Biol* 14:153-165. doi:10.1038/nrm3525
293. Venables JP, Tazi J, Juge F (2012) Regulated functional alternative splicing in *Drosophila*. *Nucleic Acids Res* 40:1-10. doi:10.1093/nar/gkr648
294. Kelemen O, Convertini P, Zhang Z, Wen Y, Shen M, *et al.* (2013) Function of alternative splicing. *Gene* 514:1-30. doi:10.1016/j.gene.2012.07.083
295. Kalsotra A, Cooper TA (2011) Functional consequences of developmentally regulated alternative splicing. *Nat Rev Genet* 12:715-729. doi:10.1038/nrg3052
296. Norris AD, Calarco JA (2012) Emerging roles of alternative pre-mRNA splicing regulation in neuronal development and function. *Front Neurosci* 6:122. doi:10.3389/fnins.2012.00122
297. Chepelev I, Chen X (2013) Alternative splicing switching in stem cell lineages. *Front Biol* 8:50-59. doi:10.1007/s11515-012-1198-y
298. Wu CY, Tsai YP, Wu MZ, Teng SC, Wu KJ (2012) Epigenetic reprogramming and post-transcriptional regulation during the epithelial-mesenchymal transition. *Trends Genet* 28:454-463. doi:10.1016/j.tig.2012.05.005
299. Warzecha CC, Carstens RP (2012) Complex changes in alternative pre-mRNA splicing play a central role in the epithelial-to-mesenchymal transition (EMT). *Semin Cancer Biol* 22:417-427. doi:10.1016/j.semcancer.2012.04.003
300. Mittendorf KE, Deatherage CL, Ohl MD, Sanders CR (2012) Tailoring of membrane proteins by alternative splicing of pre-mRNA. *Biochemistry* 51:5541-5556. doi:10.1021/bi3007065
301. Lipscombe D, Andrade A, Allen SE (2013) Alternative splicing: Functional diversity among voltage-gated calcium channels and behavioral consequences. *Biochim Biophys Acta* 1828:1522-1529. doi:10.1016/j.bbame.2012.09.018
302. Sun W, You X, Gogol-Döring A, He H, Kise Y, *et al.* (2013) Ultra-deep profiling of alternatively spliced *Drosophila* Dscam isoforms by circularization-assisted multi-segment sequencing. *EMBO J* Jun 21, 2013. doi:10.1038/emboj.2013.144
303. Peng Z, Cheng Y, Tan BC, Kang L, Tian Z, *et al.* (2012) Comprehensive analysis of RNA-Seq data reveals extensive RNA editing in a human transcriptome. *Nat Biotechnol* 30:253-260. doi:10.1038/nbt.2122
304. Bahn JH, Lee JH, Li G, Greer C, Peng G, *et al.* (2012) Accurate identification of A-to-I RNA editing in human by transcriptome sequencing. *Genome Res* 22:142-150. doi:10.1101/gr.124107.111
305. Sakurai M, Ueda H, Yano T, Okada S, Terajima H (2014) A biochemical landscape of A-to-I RNA editing in the human brain transcriptome. *Genome Res* (January 9, 2014). doi:10.1101/gr.162537.113
306. Brennicke A, Marchfelder A, Binder S (1999) RNA editing. *FEMS Microbiol Rev* 23:297-316. doi:10.1111/j.1574-6976.1999.tb00401.x

307. Eisenberg E, Li JB, Levanon EY (2010) Sequence based identification of RNA editing sites. *RNA Biol* 7:248-252. doi:10.4161/rna.7.2.11565
308. Wulff B-E, Nishikura K (2010) Substitutional A-to-I RNA editing. *Wiley Interdiscip Rev RNA* 1:90-101. doi:10.1002/wrna.10
309. Garrett S, Rosenthal JJC (2012) RNA editing underlies temperature adaptation in K⁺ channels from polar octopuses. *Science* 335:848-851. doi:10.1126/science.1212795
310. Savva YA, Rieder LE, Reenan RA (2012) The ADAR protein family. *Genome Biol* 13(12):252. doi:10.1186/gb-2012-13-12-252
311. Hoopengardner B, Bhalla T, Staber C, Reenan R (2003) Nervous system targets of RNA editing identified by comparative genomics. *Science* 301:832-836. doi:10.1126/science.1086763
312. Wahlstedt H, Daniel C, Enserö M, Ohman M (2009) Large-scale mRNA sequencing determines global regulation of RNA editing during brain development. *Genome Res* 19:978-986. doi:10.1101/gr.089409.108
313. Penn AC, Balik A, Greger IH (2013) Reciprocal regulation of A-to-I RNA editing and the vertebrate nervous system. *Front Neurosci* 7:61.
314. Li X, Overton IM, Baines RA, Keegan LP, O'Connell MA (2013) The ADAR RNA editing enzyme controls neuronal excitability in *Drosophila melanogaster*. *Nucleic Acids Res* (Epub Oct 16 2013). doi:10.1093/nar/gkt909
315. XuFeng R, Boyer MJ, Shen H, Li Y, Yu H, *et al.* (2009) ADAR1 is required for hematopoietic progenitor cell survival via RNA editing. *Proc Natl Acad Sci USA* 106:17763-17768. doi:10.1073/pnas.0903324106
316. Stulić M, Jantsch MF (2013) Spatio-temporal profiling of Filamin A RNA-editing reveals ADAR preferences and high editing levels outside neuronal tissues. *RNA Biol* 10:1611-1617. doi:10.4161/rna.26216
317. Hirata R, Ohsumi Y, Nakano A, Kawasaki H, Suzuki K, *et al.* (1990) Molecular structure of a gene, VMA1, encoding the catalytic subunit of H(+)-translocating adenosine triphosphatase from vacuolar membranes of *Saccharomyces cerevisiae*. *J Biol Chem* 265:6726-6733.
318. Kane PM, Yamashiro CT, Wolczyk DF, Neff N, Goebel M, *et al.* (1990) Protein splicing converts the yeast TFP1 gene product to the 69-kD subunit of the vacuolar H(+)-adenosine triphosphatase. *Science* 250:651-657. doi:10.1126/science.2146742
319. Paulus H (2000) Protein splicing and related forms of protein autoprocesing. *Annu Rev Biochem* 69:447-496. doi:10.1146/annurev.biochem.69.1.447
320. Perler FB (2005) Protein splicing mechanisms and applications. *IUBMB Life* 57:469-476. doi:10.1080/15216540500163343
321. Starokadomskyy PL (2007) Protein splicing. *Mol Biol* 41:278-293. doi:10.1134/S0026893307020094
322. Anraku Y, Satow Y (2009) Reflections on protein splicing: Structures, functions and mechanisms. *Proc Jpn Acad Ser B Phys Biol Sci* 85(9):409-421. doi:10.2183/pjab.85.409
323. Elleuche S, Pöggeler S (2010) Inteins, valuable genetic elements in molecular biology and biotechnology. *Appl Microbiol Biotechnol* 87:479-489. doi:10.1007/s00253-010-2628-x
324. Lee JJ, Ekker SC, von Kessler DP, Porter JA, Sun BI, *et al.* (1994) Autoproteolysis in hedgehog protein biogenesis. *Science* 266:1528-1537. doi:10.1126/science.7985023
325. Perler FB, Xu MQ, Paulus H (1997) Protein splicing and autoproteolysis mechanisms. *Curr Opin Chem Biol* 1:292-299. doi:10.1016/S1367-5931(97)80065-8
326. Rosenblum JS, Blobel G (1999) Autoproteolysis in nucleoporin biogenesis. *Proc Natl Acad Sci USA* 96:11370-11375. doi:10.1073/pnas.96.20.11370
327. Bürglin TR (2008) The hedgehog protein family. *Genome Biol* 9(11):241. doi:10.1186/gb-2008-9-11-241
328. Volkmann G, Mootz HD (2013) Recent progress in intein research: From mechanism to directed evolution and applications. *Cell Mol Life Sci* 70:1185-1206. doi:10.1007/s00018-012-1120-4
329. Dobson CM (2003) Protein folding and misfolding. *Nature* 426:884-890. doi:10.1038/nature02261
330. Sasahara K, McPhie P, Minton AP (2003) Effect of dextran on protein stability and conformation attributed to macromolecular crowding. *J Mol Biol* 326:1227-1237. doi:10.1016/S0022-2836(02)01443-2
331. Homouz D, Perham M, Samiotakis A, Cheung MS, Wittung-Stafshede P (2008) Crowded, cell-like environment induces shape changes in aspherical protein. *Proc Natl Acad Sci USA* 105:11754-11759. doi:10.1073/pnas.0803672105
332. Dhar A, Samiotakis A, Ebbinghaus S, Nienhaus L, Homouz D, *et al.* (2010) Structure, function, and folding of phosphoglycerate kinase are strongly perturbed by macromolecular crowding. *Proc Natl Acad Sci USA* 107:17586-17591. doi:10.1073/pnas.1006760107
333. Gershenson A, Gierasch LM (2011) Protein folding in the cell: Challenges and progress. *Curr Opin Struct Biol* 21:32-41. doi:10.1016/j.sbi.2010.11.001
334. Zhou HX (2013) Influence of crowded cellular environments on protein folding, binding, and oligomerization: Biological consequences and potentials of atomistic modeling. *FEBS Lett* 587:1053-1061.
335. Priestle JP, Schär H-P, Grütter MG (1988) Crystal structure of the cytokine interleukin-1 β . *EMBO J* 7:339-343.
336. Russell RB, Barton GJ (1994) Structural features can be unconserved in proteins with similar folds. *J Mol Biol* 244:332-350. doi:10.1006/jmbi.1994.1733
337. Zhang B, Jaroszewski L, Rychlewski L, Godzik A (1997) Similarities and differences between nonhomologous proteins with similar folds: Evaluation of threading strategies. *Fold Des* 2:307-317. doi:10.1016/S1359-0278(97)00042-4
338. Petros AM, Olejniczak ET, Fesik SW (2004) Structural biology of the Bcl-2 family of proteins. *Biochim Biophys Acta* 1644:83-94. doi:10.1016/j.bbamcr.2003.08.012
339. Wang C, Wu H, Katritch V, Han GW, Huang X-P, *et al.* (2013) Structure of the human smoothed receptor bound to an antitumour agent. *Nature* 497:338-343. doi:10.1038/nature12167
340. Prusiner SB (1982) Novel proteinaceous infectious particles cause scrapie. *Science* 216:136-144. doi:10.1126/science.6801762
341. Chiti F, Dobson CM (2006) Protein misfolding, functional amyloid, and human disease. *Annu Rev Biochem* 75:333-366. doi:10.1146/annurev.biochem.75.101304.123901
342. Pezza JA, Serio TR (2007) Prion propagation: The role of protein dynamics. *Prion* 1:36-43. doi:10.4161/pri.1.1.3992
343. Biljan I, Ilc G, Giachin G, Legname G, Plavec J (2013) NMR structural studies of human cellular prion proteins. *Curr Top Med Chem* 13:2407-2418. doi:10.2174/15680266113136660169
344. Murzin AG (2008) Metamorphic proteins. *Science* 320:1725-1726. doi:10.1126/science.1158868
345. Bryan PN, Orban J (2010) Proteins that switch folds. *Curr Opin Struct Biol* 20:482-488. doi:10.1016/j.sbi.2010.06.002
346. Skinner JJ, Wood S, Shorter J, Englander SW, Black BE (2008) The Mad2 partial unfolding model: Regulating mitosis through Mad2 conformational switching. *J Cell Biol* 183:761-768. doi:10.1083/jcb.200808122
347. Luo X, Yu H (2008) Protein metamorphosis: The two-state behavior of Mad2. *Structure* 16:1616-1625. doi:10.1016/j.str.2008.10.002
348. Tuinstra RL, Peterson FC, Kutlesa S, Elgin ES, Kron MA, *et al.* (2008) Interconversion between two unrelated protein folds in the lymphotactin native state. *Proc Natl Acad Sci USA* 105:5057-5062. doi:10.1073/pnas.0709518105
349. Tyler RC, Murray NJ, Peterson FC, Volkman BF (2011) Native-state interconversion of a metamorphic protein requires global unfolding. *Biochemistry* 50:7077-7079. doi:10.1021/bi200750k

350. Goodchild SC, Howell MW, Littler DR, Mandyam RA, Sale KL, *et al.* (2010) Metamorphic response of the CLIC1 chloride intracellular ion channel protein upon membrane interaction. *Biochemistry* 49:5278-5289. doi:10.1021/bi100111c
351. Valenzuela SM, Alkhamici H, Brown LJ, Almond OC, Goodchild SC, *et al.* (2013) Regulation of the membrane insertion and conductance activity of the metamorphic chloride intracellular channel protein CLIC1 by cholesterol. *PLoS One* 8(2):e56948. doi:10.1371/journal.pone.0056948
352. Dai Z, Tonelli M, Markley JL (2012) Metamorphic protein IscU changes conformation by cis-trans isomerizations of two peptidyl-prolyl peptide bonds. *Biochemistry* 51:9595-9602. doi:10.1021/bi301413y
353. Markley JL, Kim JH, Dai Z, Bothe JR, Cai K, *et al.* (2013) Metamorphic protein IscU alternates conformations in the course of its role as the scaffold protein for iron-sulfur cluster biosynthesis and delivery. *FEBS Lett* 587:1172-1179.
354. Hart FU, Bracher A, Hayer-Hart M (2011) Molecular chaperones in protein folding and proteostasis. *Nature* 475:324-332. doi:10.1038/nature10317
355. Uversky VN, Dunker AK (2010) Understanding protein non-folding. *Biochim Biophys Acta* 1804:1231-1264. doi:10.1016/j.bbapap.2010.01.017
356. Chouard T (2011) Breaking the protein rules. *Nature* 471:151-153. doi:10.1038/471151a
357. Babu MM, Kriwacki RW, Pappu RV (2012) Versatility from protein disorder. *Science* 337:1460-1461. doi:10.1126/science.1228775
358. Fuxreiter M, Tompa P (2012) Fuzzy complexes: A more stochastic view of protein function. *Adv Exp Med Biol* 725:1-14. doi:10.1007/978-1-4614-0659-4_1
359. Bozoky Z, Krzeminski M, Chong PA, Forman-Kay JD (2013) Structural changes of CFTR R region upon phosphorylation: A plastic platform for intramolecular and intermolecular interactions. *FEBS J* 280:4407-4416. doi:10.1111/febs.12422
360. Ferreon ACM, Ferreon JC, Wright PE, Deniz AA (2013) Modulation of allostery by protein intrinsic disorder. *Nature* 498:390-394. doi:10.1038/nature12294
361. Cumberworth A, Lamour G, Babu MM, Gsponer J (2013) Promiscuity as a functional trait: Intrinsically disordered regions as central players of interactomes. *Biochem J* 454:361-369. doi:10.1042/BJ20130545
362. Shental-Bechor D, Levy Y (2009) Folding of glycoproteins: Toward understanding the biophysics of the glycosylation code. *Curr Opin Struct Biol* 19:524-533. doi:10.1016/j.sbi.2009.07.002
363. Zuber C, Roth J (2009) N-glycosylation. In: Gabius H-J, ed. *The Sugar Code: Fundamentals of Glycosciences*. Wiley-VCH Verlag (Weinheim, Germany) pp 87-110.
364. Patsos G, Corfield A (2009) O-glycosylation: Structural diversity and functions. In: Gabius H-J, ed. *The Sugar Code: Fundamentals of Glycosciences*. Wiley-VCH Verlag (Weinheim, Germany) pp 111-137.
365. Aebi M (2013) N-linked protein glycosylation in the ER. *Biochim Biophys Acta* 1833:2430-2437. doi:10.1016/j.bbamcr.2013.04.001
366. Loibl M, Strahl S (2013) Protein O-mannosylation: What we have learned from baker's yeast. *Biochim Biophys Acta* 1833:2438-2446. doi:10.1016/j.bbamcr.2013.02.008
367. Potter BA, Hughey RP, Weisz OA (2006) Role of N- and O-glycans in polarized biosynthetic sorting. *Am J Physiol Cell Physiol* 290:C1-C10. doi:10.1152/ajpcell.00333.2005
368. Cullen PJ (2011) Post-translational regulation of signaling mucins. *Curr Opin Struct Biol* 21:590-596. doi:10.1016/j.sbi.2011.08.007
369. Helenius A, Aebi M (2001) Intracellular functions of N-linked glycans. *Science* 291:2364-2369. doi:10.1126/science.291.5512.2364
370. Ten Hagen KG, Bedi GS, Tetaert D, Kingsley PD, Hagen FK, *et al.* (2001) Cloning and characterization of a ninth member of the UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase family, ppGalNTase-T9. *J Biol Chem* 276:17395-17404. doi:10.1074/jbc.M009638200
371. Esko JD, Lindahl U (2001) Molecular diversity of heparan sulfate. *J Clin Invest* 108:169-173.
372. Rudd TR, Guimond SE, Skidmore MA, Duchesne L, Guerrini M, *et al.* (2007) Influence of substitution pattern and cation binding on conformation and activity in heparin derivatives. *Glycobiology* 17:983-993. doi:10.1093/glycob/cwm062
373. Hurtado-Guerrero R, Davies GJ (2012) Recent structural and mechanistic insights into post-translational enzymatic glycosylation. *Curr Opin Chem Biol* 16:479-487. doi:10.1016/j.cbpa.2012.10.013
374. Banerjee DK (2012) N-glycans in cell survival and death: Cross-talk between glycosyltransferases. *Biochim Biophys Acta* 1820:1338-1346. doi:10.1016/j.bbagen.2012.01.013
375. Bennett EP, Mandel U, Clausen H, Gerken TA, Fritz TA, *et al.* (2012) Control of mucin-type O-glycosylation: A classification of the polypeptide GalNAc-transferase gene family. *Glycobiology* 22:736-756. doi:10.1093/glycob/cwr182
376. Moremen KW, Tiemeyer M, Nairn AV (2012) Vertebrate protein glycosylation: Diversity, synthesis and function. *Nat Rev Mol Cell Biol* 13:448-462. doi:10.1038/nrm3383
377. Spiro RG (2002) Protein glycosylation: Nature, distribution, enzymatic formation, and disease implications of glycopeptide bonds. *Glycobiology* 12:43R-56R. doi:10.1093/glycob/12.4.43R
378. Lauc G, Vojta A, Zoldoš V (2014) Epigenetic regulation of glycosylation is the quantum mechanics of biology. *Biochim Biophys Acta* 1840:65-70. doi:10.1016/j.bbagen.2013.08.017
379. Palade GE (1983) Membrane biogenesis: An overview. *Method Enzymol* 96:xxix-lv. doi:10.1016/S0076-6879(83)96004-4
380. Sonneborn TM (1963) Does preformed cell structure play a role in heredity? In: Allen JM, ed. *The Nature of Biological Diversity*. McGraw-Hill (New York) pp 165-221.
381. Beisson J, Sonneborn TM (1965) Cytoplasmic inheritance of the organization of the cell cortex in *Paramecium aurelia*. *Proc Natl Acad Sci USA* 53:275-282. doi:10.1073/pnas.53.2.275
382. Grimes GW, Aufderheide KJ (1991) Cellular Aspects of Pattern Formation: The Problem of Assembly. Karger (Basel) pp 22-26.
383. Ng SF, Frankel J (1977) 180° rotation of ciliary rows and its morphogenetic implications in *Tetrahymena pyriformis*. *Proc Natl Acad Sci USA* 74:1115-1118. doi:10.1073/pnas.74.3.1115
384. Frankel J (1980) Propagation of cortical differences in *Tetrahymena*. *Genetics* 94:607-623.
385. Nanney DL (1968) Cortical patterns in cellular morphogenesis. *Science* 160:496-502. doi:10.1126/science.160.3827.496
386. Grimes GW (1982) Pattern determination in hypotrich ciliates. *Am Zool* 22:35-46.
387. Sonneborn TM (1930) Genetic studies in *Stenostomum incaudatum* (Nov. spec.). I. The nature and origin of differences among individuals formed during vegetative reproduction. *J Exp Zool* 57:57-108. doi:10.1002/jez.1400570104
388. Sonneborn TM (1930) Genetic studies in *Stenostomum incaudatum*. II. The effects of lead acetate on the hereditary constitution. *J Exp Zool* 57:409-439. doi:10.1002/jez.1400570303
389. Locke M (1990) Is there somatic inheritance of intracellular patterns? *J Cell Sci* 96:563-567.
390. Poyton RO (1983) Memory and membranes: The expression of genetic and spatial memory during the assembly of organelle macrocompartments. *Mod Cell Biol* 2:15-72.
391. Lee AG (2004) How lipids affect the activities of integral membrane proteins. *Biochim Biophys Acta* 1666:62-87. doi:10.1016/j.bbamem.2004.05.012
392. Li E, Wimley WC, Hristova K (2012) Transmembrane helix dimerization: Beyond the search for sequence motifs. *Biochim Biophys Acta* 1818:183-193. doi:10.1016/j.bbamem.2011.08.031
393. Cymer F, Veerappan A, Schneider D (2012) Transmembrane helix-helix interactions are modulated by the sequence context and by lipid bilayer properties. *Biochim Biophys Acta* 1818:963-973. doi:10.1016/j.bbamem.2011.07.035

394. Cristian L, Lear JD, DeGrado WF (2003) Use of thiol-disulfide equilibria to measure the energetics of assembly of transmembrane helices in phospholipid bilayers. *Proc Natl Acad Sci USA* 100:14772-14777. doi:10.1073/pnas.2536751100
395. You M, Li E, Wimley WC, Hristova K (2005) Förster resonance energy transfer in liposomes: Measurements of transmembrane helix dimerization in the native bilayer environment. *Anal Biochem* 340:154-164. doi:10.1016/j.ab.2005.01.035
396. Katzmann DJ, Stefan CJ, Babst M, Emr SD (2003) Vps27 recruits ESCRT machinery to endosomes during MVB sorting. *J Cell Biol* 162:413-423. doi:10.1083/jcb.200302136
397. Hong H, Blois TM, Cao Z, Bowie JU (2010) Method to measure strong protein-protein interactions in lipid bilayers using a steric trap. *Proc Natl Acad Sci USA* 107:19802-19807. doi:10.1073/pnas.1010348107
398. Fyfe I, Schuh AL, Edwardson JM, Audhya A (2011) Association of the endosomal sorting complex ESCRT-II with the Vps20 subunit of ESCRT-III generates a curvature-sensitive complex capable of nucleating ESCRT-III filaments. *J Biol Chem* 286:34262-34270. doi:10.1074/jbc.M111.266411
399. Adell MA, Teis D (2011) Assembly and disassembly of the ESCRT-III membrane scission complex. *FEBS Lett* 585:3191-3196.
400. Plowman SJ, Ariotti N, Goodall A, Parton RG, Hancock JF (2008) Electrostatic interactions positively regulate K-Ras nanocluster formation and function. *Mol Cell Biol* 28:4377-4385. doi:10.1128/MCB.00050-08
401. Abankwa D, Gorfieb AA, Indera K, Hancock JF (2010) Ras membrane orientation and nanodomain localization generate isoform diversity. *Proc Natl Acad Sci USA* 107:1130-1135. doi:10.1073/pnas.0903907107
402. Menendez-Benito V, van Deventer SJ, Jimenez-Garcia V, Roy-Luzarraga M, van Leeuwen F, et al. (2013) Spatiotemporal analysis of organelle and macromolecular complex inheritance. *Proc Natl Acad Sci USA* 110:175-80. doi:10.1073/pnas.1207424110
403. Cavalier-Smith T (2004) The membranome and membrane heredity in development and evolution. In: Hirt RP, Horner DS, eds. *Organelles, Genomes and Eukaryote Phylogeny*. CRC Press (Boca Raton, FL) pp 335-351.
404. Ghosh D, Beavis RC, Wilkins JA (2008) The identification and characterization of membranome components. *J Proteome Res* 7:1572-1583. doi:10.1021/pr070509u
405. Li Y, Yu J, Wang Y, Griffin NM, Long F, et al. (2009) Enhancing identifications of lipid-embedded proteins by mass spectrometry for improved mapping of endothelial plasma membranes in vivo. *Mol Cell Proteomics* 8:1219-1235. doi:10.1074/mcp.M800215-MCP200
406. Uva P, Lahm A, Sbardellati A, Grigoriadis A, Tutt A, et al. (2010) Comparative membranome expression analysis in primary tumors and derived cell lines. *PLoS One* 5(7):e11742. doi:10.1371/journal.pone.0011742
407. Cavalier-Smith T (2000) Membrane heredity and early chloroplast evolution. *Trends Plant Sci* 5:174-182. doi:10.1016/S1360-1385(00)01598-3
408. Tatewaki M, Nagata K (1970) Surviving protoplasts *in vitro* and their development in *Bryopsis*. *J Phycol* 6:401-403. doi:10.1111/j.1529-8817.1970.tb02414.x
409. Kobayashi K, Kanaizuka Y (1985) Reunification of sub-cellular fractions of *Bryopsis* into viable cells. *Plant Sci* 40:129-135. doi:10.1016/0168-9452(85)90053-6
410. Pak JY, Solorzano C, Arai M, Nitta T (1991). Two distinct steps for spontaneous generation of subprotoplasts from a disintegrated *Bryopsis* cell. *Plant Physiol* 96:819-825.
411. Kim GH, Klotchkova TA, Kang YM (2001) Life without a cell membrane: Regeneration of protoplasts from disintegrated cells of the marine green alga *Bryopsis plumosa*. *J Cell Sci* 114:2009-2014.
412. Wang G-C, Tseng C-K (2006) Culturing the segments of *Bryopsis hypnoides* Lamouroux Thalli regenerated from protoplast aggregations. *J Integr Plant Biol* 48:190-196. doi:10.1111/j.1744-7909.2006.00190.x
413. Ram M, Babbar SB (2002) Transient existence of life without a cell membrane: A novel strategy of siphonous seaweed for survival and propagation. *BioEssays* 24:588-590. doi:10.1002/bies.10117
414. Wells J (2013) The membrane code: A carrier of essential biological information that is not specified by DNA and is inherited apart from it. In: Marks RJ II, Behe MJ, Dembski WA, Gordon BL, Sanford JC, eds. *Biological Information: New Perspectives*. World Scientific (Singapore) pp 474-488.
415. Darwin C (1859) *On the Origin of Species by Means of Natural Selection*. John Murray (London) p 131.
416. Butler S (1880) *Unconscious Memory*. David Bogue (London) ch 13.
417. Romanes GJ (1895) *Darwin, and After Darwin*. Longmans, Green (London) v 2 pp 12-13.
418. Huxley JS (1942) *Evolution: The Modern Synthesis*. Allen & Unwin (London).
419. Mayr E (1982) *The Growth of Biological Thought*. Harvard University Press (Cambridge, MA), pp 537,566-570.
420. Gould SJ (2002) *The Structure of Evolutionary Theory*. Harvard University Press (Cambridge, MA) p 216.
421. Haldane JBS (1927) A mathematical theory of natural and artificial selection, part V: Selection and mutation. *Math Proc Cambridge* 23:838-844. doi:10.1017/S0305004100015644
422. Fisher RA (1930) *The Genetical Theory of Natural Selection*. Clarendon Press (Oxford).
423. Wright SG (1931) Evolution in Mendelian populations. *Genetics* 16:97-159.
424. Lynch M (2007) *The Origins of Genome Architecture*. Sinauer Associates (Sunderland, MA) p 370.
425. Hall BK (1992) *Evolutionary Developmental Biology*. Chapman & Hall (London) pp ix,9,223.
426. Carroll SB (2005) *Endless Forms Most Beautiful: The New Science of Evo-Devo*. W. W. Norton (New York) pp 35,90-91,116.
427. Tyson JJ (1976) The Belousov-Zhabotinsky Reaction. In: Levin S, ed. *Lecture Notes in Mathematics*. (Springer-Verlag (Berlin) v10.
428. Pechenkin A (2009) B. P. Belousov and his reaction. *J Biosci* 34:365-371. doi:10.1007/s12038-009-0042-2
429. Bedau M (1997) Weak emergence. In: Tomberlin J, ed. *Philosophical Perspectives: Mind, Causation, and World*. Blackwell (Malden, MA) v2, pp 375-399.
430. Kauffman SA (1995) *At Home in the Universe: The Search for Laws of Self-Organization and Complexity*. Oxford University Press (New York).
431. Chalmers DJ (2006) Strong and weak emergence. In: Clayton P, Davies P, eds. *The Re-emergence of Emergence*. Oxford University Press (Oxford).
432. Rashevsky N (1954) Topology and life: In search of general mathematical principles in biology and sociology. *Bull Math Biophys* 16:317-348. doi:10.1007/BF02484495
433. Rosen R (1958) A relational theory of biological systems. *Bull Math Biophys* 20:245-260. doi:10.1007/BF02478302
434. Rashevsky N (1960) Contributions to relational biology. *Bull Math Biophys* 22:73-84. doi:10.1007/BF02477973
435. Rosen R (1991) *Life Itself: A Comprehensive Inquiry Into the Nature, Origin, and Fabrication of Life*. Columbia University Press (New York).
436. Eilenberg S, Mac Lane S (1945) General theory of natural equivalences. *Trans Amer Math Soc* 58:231-294. doi:10.1090/S0002-9947-1945-0013131-6
437. Grothendieck A (1957) Sur quelques points d'algèbre homologique. *Tôhoku Math J* 9:199-221.

438. Freyd PJ (1964) *Abelian Categories: An Introduction to the Theory of Functors*. Harper & Row (New York).
439. Lawvere FW (1963) Functorial semantics of algebraic theories. *Proc Natl Acad Sci USA* 50:869-872. doi:10.1073/pnas.50.5.869
440. Lawvere FW (1964) An elementary theory of the category of sets. *Proc Natl Acad Sci USA* 52:1506-1511. doi:10.1073/pnas.52.6.1506
441. Awodey S (2010) *Category Theory*. Second Edition. Oxford University Press (Oxford).
442. MacLane S (1998) *Categories for the Working Mathematician*. Second Edition. Springer-Verlag (New York).
443. Lawvere FW, Schanuel SH (2009) *Conceptual Mathematics: A First Introduction to Categories*. Second Edition. Cambridge University Press (Cambridge).
444. Rosen R (1958) A representation of biological systems from the standpoint of the theory of categories. *Bull Math Biophys* 20:317-341. doi:10.1007/BF02477890
445. Baianu I (1980) Natural transformations of organismic structures. *Bull Math Biol* 42:431-446. doi:10.1007/BF02460794
446. Ehresmann AC, Vanbremeersch J-P (2007) *Memory Evolutive Systems*. Studies in Multidisciplinarity v 4. Elsevier (Amsterdam).
447. Sternberg R (2008) DNA codes and information: Formal structures and relational causes. *Acta Biotheor* 56:205-232. doi:10.1007/s10441-008-9049-6
448. Louie AH (2009) *More Than Life Itself: A Synthetic Continuation in Relational Biology*. Ontos Verlag (Frankfurt).